

Research Article

Existence and Data Dependence of Fixed Points and Strict Fixed Points for Contractive-Type Multivalued Operators

Cristian Chifu and Gabriela Petrușel

Received 21 October 2006; Revised 1 December 2006; Accepted 2 December 2006

Recommended by Simeon Reich

The purpose of this paper is to present several existence and data dependence results of the fixed points of some multivalued generalized contractions in complete metric spaces. As for application, a continuation result is given.

Copyright © 2007 C. Chifu and G. Petrușel. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Throughout this paper, the standard notations and terminologies in nonlinear analysis (see [14, 15]) are used. For the convenience of the reader we recall some of them.

Let (X, d) be a metric space. By $\tilde{B}(x_0, r)$ we denote the closed ball centered in $x_0 \in X$ with radius $r > 0$.

Also, we will use the following symbols:

$$\begin{aligned} P(X) &:= \{Y \subset X \mid Y \text{ is nonempty}\}, & P_{\text{cl}}(X) &:= \{Y \in P(X) \mid Y \text{ is closed}\}, \\ P_b(X) &:= \{Y \in P(X) \mid Y \text{ is bounded}\}, & P_{b,\text{cl}}(X) &:= P_{\text{cl}}(X) \cap P_b(X). \end{aligned} \quad (1.1)$$

Let A and B be nonempty subsets of the metric space (X, d) . The gap between these sets is

$$D(A, B) = \inf \{d(a, b) \mid a \in A, b \in B\}. \quad (1.2)$$

In particular, $D(x_0, B) = D(\{x_0\}, B)$ (where $x_0 \in X$) is called the distance from the point x_0 to the set B .

2 Fixed Point Theory and Applications

The Pompeiu-Hausdorff generalized distance between the nonempty closed subsets A and B of the metric space (X, d) is defined by the following formula:

$$H(A, B) := \max \left\{ \sup_{a \in A} \inf_{b \in B} d(a, b), \sup_{b \in B} \inf_{a \in A} d(a, b) \right\}. \quad (1.3)$$

If $A, B \in P_{b,cl}(X)$, then one denotes

$$\delta(A, B) := \sup \{d(a, b) \mid a \in A, b \in B\}. \quad (1.4)$$

The symbol $T : X \rightarrow P(Y)$ denotes a set-valued operator from X to Y . We will denote by $\text{Graph}(T) := \{(x, y) \in X \times Y \mid y \in T(x)\}$ the graph of T . Recall that the set-valued operator is called closed if $\text{Graph}(T)$ is a closed subset of $X \times Y$.

For $T : X \rightarrow P(X)$ the symbol $\text{Fix}(T) := \{x \in X \mid x \in T(x)\}$ denotes the fixed point set of the set-valued operator T , while $\text{SFix}(T) := \{x \in X \mid \{x\} = T(x)\}$ is the strict fixed point set of T .

If (X, d) is a metric space, $T : X \rightarrow P_{cl}(X)$ is called a multivalued a -contraction if $a \in]0, 1[$ and $H(T(x_1), T(x_2)) \leq a \cdot d(x_1, x_2)$, for each $x_1, x_2 \in X$.

In the same setting, an operator $T : X \rightarrow P_{cl}(X)$ is a multivalued weakly Picard operator (briefly MWP operator) (see [15]) if for each $x \in X$ and each $y \in T(x)$ there exists a sequence $(x_n)_{n \in \mathbb{N}}$ in X such that

- (i) $x_0 = x, x_1 = y$,
- (ii) $x_{n+1} \in T(x_n)$, for all $n \in \mathbb{N}$,
- (iii) the sequence $(x_n)_{n \in \mathbb{N}}$ is convergent and its limit is a fixed point of T .

Any multivalued a -contraction or any multivalued Reich-type operator (see Reich [10]) are examples of MWP operators. For other examples and results, see Petruşel [9].

Also, let us mention that a sequence $(x_n)_{n \in \mathbb{N}}$ in X satisfying the condition (ii) from the previous definition is called the sequence of successive approximations of T starting from $x_0 \in X$.

The following result was proved in the work of Feng and Liu (see [5]).

THEOREM 1.1 (Feng, Liu). *Let (X, d) be a complete metric space, $T : X \rightarrow P_{cl}(X)$ and $q > 1$. Consider $S_q(x) := \{y \in T(x) \mid d(x, y) \leq q \cdot D(x, T(x))\}$. Suppose that T satisfies the following condition:*

- (1.1) *there is a $a < 1/q$ such that for each $x \in X$ there is $y \in S_q(x)$ satisfying*

$$D(y, T(y)) \leq a \cdot d(x, y). \quad (1.5)$$

Also, suppose that the function $p : X \rightarrow \mathbb{R}$, $p(x) := D(x, T(x))$ is lower semicontinuous. Then $\text{Fix } T \neq \emptyset$.

The purpose of this paper is to study the existence and data dependence of the fixed points and strict fixed points for some self and nonself multivalued operators satisfying to some generalized Feng-Liu-type conditions.

Our results are in connection with the theory of MWP operators (see [9, 15]) and they generalize some fixed point and strict fixed point principles for multivalued operators given in [3–5, 7, 8, 10–13].

2. Fixed points

Let (X, d) be a metric space, $T : X \rightarrow P_{cl}(X)$ a multivalued operator, and $q > 1$. Define $S_q(x) := \{y \in T(x) \mid d(x, y) \leq q \cdot D(x, T(x))\}$. Obviously $S_q(x) \neq \emptyset$, for each $x \in X$ and S_q is a multivalued selection of T .

Our first main result is the following theorem.

THEOREM 2.1. *Let (X, d) be a complete metric space, $x_0 \in X$, $r > 0$, $q > 1$, and $T : X \rightarrow P_{cl}(X)$ a multivalued operator. Suppose that*

- (i) *there exists $a \in \mathbb{R}_+$ with $aq < 1$ such that for each $x \in \tilde{B}(x_0, r)$ there exists $y \in S_q(x)$ having the property*

$$D(y, T(y)) \leq a \cdot d(x, y), \quad (2.1)$$

- (ii) *T is closed or the function $p : X \rightarrow \mathbb{R}_+$, $p(x) := D(x, T(x))$ is lower semicontinuous,*
 (iii) *$D(x_0, T(x_0)) \leq ((1 - aq)/q) \cdot r$.*

Then $\text{Fix}(T) \cap \tilde{B}(x_0, r) \neq \emptyset$.

Proof. From (i) and (iii) there is $x_1 \in T(x_0)$ such that $d(x_0, x_1) \leq qD(x_0, T(x_0)) < (1 - aq)r$ and $D(x_1, T(x_1)) \leq ad(x_0, x_1) \leq aqD(x_0, T(x_0))$. Hence $x_1 \in \tilde{B}(x_0, r)$. Next, we can find $x_2 \in T(x_1)$ such that $d(x_1, x_2) \leq qD(x_1, T(x_1)) \leq aqd(x_0, x_1) < aq(1 - aq) \cdot r$ and $D(x_2, T(x_2)) \leq ad(x_1, x_2) \leq aqD(x_1, T(x_1)) \leq (aq)^2D(x_0, T(x_0))$. As a consequence, $d(x_0, x_2) \leq d(x_0, x_1) + d(x_1, x_2) \leq (1 - aq)r + aq(1 - aq)r = (1 - (aq)^2)r$ and so $x_2 \in \tilde{B}(x_0, r)$.

Inductively we get a sequence $(x_n)_{n \in \mathbb{N}}$ having the following properties:

- (a) $x_{n+1} \in T(x_n)$, $n \in \mathbb{N}$;
 (b) $d(x_n, x_{n+1}) \leq (aq)^n d(x_0, x_1)$, $d(x_0, x_n) \leq (1 - (aq)^n)r$, $n \in \mathbb{N}$;
 (c) $D(x_n, T(x_n)) \leq (aq)^n \cdot D(x_0, T(x_0))$, $n \in \mathbb{N}$.

From (b) we have that $(x_n)_{n \in \mathbb{N}}$ converges to $x^* \in \tilde{B}(x_0, r)$.

From (a) and the fact that $\text{Graph}T$ is closed we obtain $x^* \in \text{Fix}T$.

From (c) and the fact that p is lower semicontinuous we have $p(x_n) \leq (aq)^n \cdot p(x_0)$, for each $n \in \mathbb{N}$. Since $aq < 1$, we immediately deduce that the sequence $(p(x_n))$ is convergent to 0, as $n \rightarrow +\infty$. Then $0 \leq p(x^*) \leq \liminf_{n \rightarrow +\infty} p(x_n) = 0$. So, $p(x^*) = 0$ and then $x^* \in T(x^*)$. \square

Remark 2.2. The above result is a local version of the main result in [5, Theorem 3.1] see Theorem 1.1. In particular, Theorem 1.1 follows from Theorem 2.1 by taking $r := +\infty$. Theorem 2.1 also extends some results from [3, 4, 7–9], and so forth.

As for application, a homotopy result can be proved.

THEOREM 2.3. *Let (X, d) be a complete metric space, U an open subset of X , and $q > 1$. Suppose that $G : \overline{U} \times [0, 1] \rightarrow P_{cl}(X)$ is a closed multivalued operator such that the following conditions are satisfied:*

- (a) *$x \notin G(x, t)$, for each $x \in \partial U$ and each $t \in [0, 1]$;*
 (b) *there exists $a \in \mathbb{R}_+$ with $aq < 1$, such that for each $t \in [0, 1]$ and each $x \in \overline{U}$ there exists $y \in \overline{U} \cap S_q(x, t)$ (where $S_q(x, t) := \{y \in G(x, t) \mid d(x, y) \leq q \cdot D(x, G(x, t))\}$),*

4 Fixed Point Theory and Applications

with the property

$$D(y, G(y, t)) \leq a \cdot d(x, y); \quad (2.2)$$

(c) there exists a continuous increasing function $\phi : [0, 1] \rightarrow \mathbb{R}$ such that

$$H(G(x, t), G(x, s)) \leq |\phi(t) - \phi(s)| \quad \forall t, s \in [0, 1] \text{ and each } x \in \bar{U}. \quad (2.3)$$

Then $G(\cdot, 0)$ has a fixed point if and only if $G(\cdot, 1)$ has a fixed point.

Proof. Suppose $G(\cdot, 0)$ has a fixed point. Define

$$Q := \{(t, x) \in [0, 1] \times U \mid x \in G(x, t)\}. \quad (2.4)$$

Obviously $Q \neq \emptyset$. Consider on Q a partial order defined as follows:

$$(t, x) \leq (s, y) \quad \text{iff } t \leq s, \quad d(x, y) \leq \frac{2q}{1-aq} \cdot [\phi(s) - \phi(t)]. \quad (2.5)$$

Let M be a totally ordered subset of Q and consider $t^* := \sup\{t \mid (t, x) \in M\}$. Consider a sequence $(t_n, x_n)_{n \in \mathbb{N}^*} \subset M$ such that $(t_n, x_n) \leq (t_{n+1}, x_{n+1})$ and $t_n \rightarrow t^*$, as $n \rightarrow +\infty$. Then

$$d(x_m, x_n) \leq \frac{2q}{1-aq} \cdot [\phi(t_m) - \phi(t_n)], \quad \text{for each } m, n \in \mathbb{N}^*, m > n. \quad (2.6)$$

When $m, n \rightarrow +\infty$ we obtain $d(x_m, x_n) \rightarrow 0$ and so $(x_n)_{n \in \mathbb{N}^*}$ is Cauchy. Denote by $x^* \in X$ its limit. Then $x_n \in G(x_n, t^*)$, $n \in \mathbb{N}^*$ and G closed imply that $x^* \in G(x^*, t^*)$. Also, from (a) $x^* \in U$. Hence $(t^*, x^*) \in Q$. Since M is totally ordered, we get $(t, x) \leq (t^*, x^*)$, for each $(t, x) \in M$. Thus (t^*, x^*) is an upper bound of M . Hence Zorn's lemma applies and Q admits a maximal element $(t_0, x_0) \in Q$. We claim that $t_0 = 1$. This will finish the proof.

Suppose the contrary, that is, $t_0 < 1$. Choose $r > 0$ and $t \in]t_0, 1]$ such that $\tilde{B}(x_0, r) \subset U$ and $r := (2q/(1-aq)) \cdot [\phi(t) - \phi(t_0)]$.

Then the $D(x_0, G(x_0, t)) \leq D(x_0, G(x_0, t_0)) + H(G(x_0, t_0), G(x_0, t)) \leq 0 + [\phi(t) - \phi(t_0)] = (1-aq)r/2q < (1-aq)r/q$.

Then the multivalued operator $G(\cdot, t) : \tilde{B}(x_0, r) \rightarrow P_{cl}(X)$ satisfies all the assumptions of Theorem 2.1. Hence there exists a fixed point $x \in \tilde{B}(x_0, r)$ for $G(\cdot, t)$. Thus $(t, x) \in Q$. Since

$$d(x_0, x) \leq r = \frac{2q}{1-aq} \cdot [\phi(t) - \phi(t_0)], \quad (2.7)$$

we immediately get $(t_0, x_0) < (t, x)$. This is a contradiction with the maximality of (t_0, x_0) . \square

Remark 2.4. Theorem 2.3 extends the main theorem in the work of Frigon and Granas [6]. See also Agarwal et al. [1] and Chiş and Precup [2] for some similar results or possibilities for extension.

Another fixed point result is the following.

THEOREM 2.5. *Let (X, d) be a complete metric space, $x_0 \in X$, $r > 0$, $q > 1$, and $T : X \rightarrow P_{cl}(X)$ a multivalued operator. Suppose that*

- (i) *there exists $a, b \in \mathbb{R}_+$ with $aq + b < 1$ such that for each $x \in \tilde{B}(x_0, r)$ there exists $y \in S_q(x)$ having the property*

$$D(y, T(y)) \leq a \cdot d(x, y) + b \cdot D(x, T(x)), \quad (2.8)$$

(ii) *T is closed or the function $p : X \rightarrow \mathbb{R}_+$, $p(x) := D(x, T(x))$ is lower semicontinuous,*

(iii) *$D(x_0, T(x_0)) < ((1 - (aq + b))/q) \cdot r$.*

Then $\text{Fix}(T) \cap \tilde{B}(x_0, r) \neq \emptyset$.

Proof. By (i) and (iii) we deduce the existence of an element $x_1 \in T(x_0)$ such that $d(x_0, x_1) \leq qD(x_0, T(x_0)) < (1 - (aq + b))r$ and $D(x_1, T(x_1)) \leq ad(x_0, x_1) + bD(x_0, T(x_0)) \leq (aq + b)D(x_0, T(x_0))$.

Inductively we obtain $(x_n)_{n \in \mathbb{N}}$ a sequence of successive approximations of T satisfying, for each $n \in \mathbb{N}$, the following relations:

- (1) $d(x_n, x_{n+1}) \leq q(aq + b)^n \cdot D(x_0, T(x_0))$, $d(x_0, x_n) \leq (1 - (aq + b)^n) \cdot r$,
(2) $D(x_n, T(x_n)) \leq (aq + b)^n \cdot D(x_0, T(x_0))$.

The rest of the proof runs as before and so the conclusion follows. \square

Remark 2.6. The above result generalizes the fixed point result in the work of Rus [12], where the following graphic contraction condition is involved: there is $a, b \in \mathbb{R}_+$ with $a + b < 1$ such that $H(T(x), T(y)) \leq a \cdot d(x, y) + bD(x, T(x))$, for each $x \in X$ and each $y \in T(x)$.

A data dependence result is the following.

THEOREM 2.7. *Let (X, d) be a complete metric space, $T_1, T_2 : X \rightarrow P_{cl}(X)$ multivalued operators, and $q_1, q_2 > 1$. Suppose that*

- (i) *there exist $a_i, b_i \in \mathbb{R}_+$ with $a_i q_i + b_i < 1$ such that for each $x \in X$ there exists $y \in S_{q_i}(x)$ having the property*

$$D(y, T_i(y)) \leq a_i \cdot d(x, y) + b_i \cdot D(x, T_i(x)), \quad \text{for } i \in \{1, 2\}; \quad (2.9)$$

(ii) *there exists $\eta > 0$ such that $H(T_1(x), T_2(x)) \leq \eta$, for each $x \in X$;*

(iii) *T_i is closed or the function $p_i : X \rightarrow \mathbb{R}_+$, $p_i(x) := D(x, T_i(x))$ is lower semicontinuous, for $i \in \{1, 2\}$.*

Then

(a) $\text{Fix}(T_i) \in P_{cl}(X)$, for $i \in \{1, 2\}$,

(b) $H(\text{Fix}(T_1), \text{Fix}(T_2)) \leq \max_{i \in \{1, 2\}} \{q_i / (1 - (a_i q_i + b_i))\} \cdot \eta$.

Proof. (a) By Theorem 2.1 we have that $\text{Fix } T_i \neq \emptyset$, for $i \in \{1, 2\}$. Also, $\text{Fix } T_i$ is closed, for $i \in \{1, 2\}$. Indeed, for example, let $(u_n)_{n \in \mathbb{N}} \in \text{Fix } T_1$, such that $u_n \rightarrow u$, as $n \rightarrow +\infty$. Then, when T_1 is closed, the conclusion follows. When $p_1(x) := D(x, T_1(x))$ is lower semicontinuous we have $0 \leq p_1(u) \leq \liminf_{n \rightarrow +\infty} p_1(u_n) = 0$. Hence $p_1(u) = 0$ and so $u \in \text{Fix } T_1$.

6 Fixed Point Theory and Applications

(b) For the second conclusion, let $x_0^* \in \text{Fix } T_1$. Then there exists $x_1 \in S_{q_2}(x_0^*)$ with $D(x_1, T_2(x_1)) \leq a_2 \cdot d(x_0^*, x_1) + b_2 \cdot D(x_0^*, T_2(x_0^*))$. Hence $d(x_0^*, x_1) \leq q_2 \cdot D(x_0^*, T_2(x_0^*))$ and $D(x_1, T_2(x_1)) \leq (a_2 q_2 + b_2) \cdot D(x_0^*, T_2(x_0^*))$. Inductively we get a sequence $(x_n)_{n \in \mathbb{N}}$ with the following properties:

- (1) $x_0 = x_0^* \in \text{Fix } T_1$,
- (2) $d(x_n, x_{n+1}) \leq q_2(a_2 q_2 + b_2)^n \cdot D(x_0^*, T_2(x_0^*))$, $n \in \mathbb{N}$,
- (3) $D(x_n, T_2(x_n)) \leq (a_2 q_2 + b_2)^n \cdot D(x_0^*, T_2(x_0^*))$, $n \in \mathbb{N}$.

From (2) we have

$$d(x_n, x_{n+m}) \leq q_2(a_2 q_2 + b_2)^n \cdot \frac{1 - (a_2 q_2 + b_2)^m}{1 - (a_2 q_2 + b_2)} D(x_0^*, T_2(x_0^*)). \quad (2.10)$$

Hence $(x_n)_{n \in \mathbb{N}}$ is Cauchy and so it converges to an element $u_2^* \in X$. As in the proof of Theorem 2.1, from (3) we immediately get that $u_2^* \in \text{Fix } T_2$. When $m \rightarrow +\infty$ in the above relation, we obtain $d(x_n, u_2^*) \leq (q_2(a_2 q_2 + b_2)^n / (1 - (a_2 q_2 + b_2))) D(x_0^*, T_2(x_0^*))$, for each $n \in \mathbb{N}$.

For $n = 0$ we get $d(x_0, u_2^*) \leq q_2 / (1 - (a_2 q_2 + b_2)) D(x_0^*, T_2(x_0^*))$.

As a consequence

$$d(x_0, u_2^*) \leq \frac{q_2}{1 - (a_2 q_2 + b_2)} \cdot H(T_1(x_0^*), T_2(x_0^*)) \leq \frac{q_2}{1 - (a_2 q_2 + b_2)} \cdot \eta. \quad (2.11)$$

In a similar way we can prove that for each $y_0^* \in \text{Fix } T_2$ there exists $u_1^* \in \text{Fix } T_1$ such that $d(y_0, u_1^*) \leq q_1 / (1 - (a_1 q_1 + b_1)) \cdot \eta$. The proof is complete. \square

Remark 2.8. Theorem 2.7 gives (for $b_i = 0$, $i \in \{1, 2\}$) a data dependence result for the fixed point set of a generalized contraction in Feng and Liu sense, see [5].

Remark 2.9. The condition $D(T(x), T(y)) \leq a \cdot d(x, y)$, for each $x, y \in X$, does not imply the existence of a fixed point for a multivalued operator $T : X \rightarrow P_{\text{cl}}(X)$. Take for example $X := [1, +\infty]$ and $T(x) := [2x, +\infty[$ see also [10]. On the other hand, if $X := \{0, 1\} \cup \{k^n \mid n \in \mathbb{N}^*\}$ (with $k \in]0, 1[$) and $T : X \rightarrow P_{\text{cl}}(X)$ given by

$$T(x) = \begin{cases} \{0, k\}, & \text{if } x = 0, \\ \{k^{n+1}, 1\}, & \text{if } x = k^n \ (n \in \mathbb{N}), \end{cases} \quad (2.12)$$

then T does not satisfies the hypothesis of Nadler's theorem, but satisfies the condition $D(y, T(y)) \leq a \cdot d(x, y) + b \cdot D(x, T(x))$, for each $(x, y) \in \text{Graph } T$ and $\text{Fix } T = \{0\}$.

3. Strict fixed points

Let (X, d) be a metric space, $T : X \rightarrow P_{b, \text{cl}}(X)$ a multivalued operator, and $q > 1$. Define $M_q(x) := \{y \in T(x) \mid \delta(x, T(x)) \leq q \cdot d(x, y)\}$. Obviously, M_q is a multivalued selection of T and $M_q(x) \neq \emptyset$, for each $x \in X$.

We have the following theorem.

THEOREM 3.1. *Let (X, d) be a complete metric space, $T : X \rightarrow P_b(X)$ a multivalued operator and $q > 1$. Suppose*

(3.1) *there exists $a \in \mathbb{R}_+$ with $aq < 1$ such that for each $x \in X$ there exists $y \in M_q(x)$ having the property*

$$\delta(y, T(y)) \leq a \cdot \max \left\{ \delta(x, T(x)), \frac{1}{2} [D(x, T(y))] \right\}. \quad (3.1)$$

If the function $r : X \rightarrow \mathbb{R}_+$, $r(x) := \delta(x, T(x))$ is lower semicontinuous, then $\text{SFix}(T) \neq \emptyset$.

Proof. Let $x_0 \in X$. If $\delta(x_0, T(x_0)) = 0$ we are done. Suppose that $\delta(x_0, T(x_0)) > 0$. Then there exists $x_1 \in M_q(x_0)$ such that $\delta(x_1, T(x_1)) \leq a \cdot \max \{ \delta(x_0, T(x_0)), (1/2) \cdot D(x_0, T(x_1)) \} \leq \max \{ a/(2-a), aq \} d(x_0, x_1)$.

Inductively we construct a sequence $(x_n)_{n \in \mathbb{N}}$ of successive approximation of T with $\delta(x_n, T(x_n)) \leq q \cdot d(x_n, x_{n+1})$, for each $n \in \mathbb{N}$. Then $d(x_n, x_{n+1}) \leq \delta(x_n, T(x_n)) \leq a \cdot \max \{ \delta(x_{n-1}, T(x_{n-1})), (1/2) \cdot D(x_{n-1}, T(x_n)) \} \leq a \cdot \max \{ q \cdot d(x_{n-1}, x_n), (1/2) \cdot D(x_{n-1}, T(x_n)) \} \leq \max \{ a/(2-a), aq \} \cdot d(x_{n-1}, x_n)$. Since $\alpha := \max \{ a/(2-a), aq \} < 1$, we immediately get that the sequence $(x_n)_{n \in \mathbb{N}}$ is convergent in the complete metric space (X, d) . Denote by x^* its limit.

We also have that $r(x_{n+1}) \leq q \cdot \alpha^n \cdot d(x_0, x_1)$. When $n \rightarrow +\infty$ we obtain $\lim_{n \rightarrow +\infty} r(x_n) = 0$. From the lower semicontinuity of r we conclude $0 \leq r(x^*) \leq \liminf_{n \rightarrow +\infty} r(x_n) = 0$. Hence $\delta(x^*, T(x^*)) = 0$ and so $x^* \in \text{SFix } T$. \square

Remark 3.2. The above result generalizes some strict fixed point results, given by Reich in [10, 11], Rus in [12, 13] and Ćirić in [3]. In particular, (3.1) implies the Ćirić-type condition on the graph of T .

Remark 3.3. If X is a metric space, the condition

$$\delta(T(x), T(y)) \leq a \cdot d(x, y), \quad \text{for each } x, y \in X, \quad (3.2)$$

necessarily implies that T is singlevalued. This is not the case, if T satisfies the condition

$$\delta(y, T(y)) \leq a \cdot \max \left\{ d(x, y), \delta(x, T(x)), \frac{1}{2} [D(x, T(y)) + D(y, T(x))] \right\}, \quad (3.3)$$

for each $(x, y) \in X$. Take for example $X := [0, 1]$ and $T(x) := [0, x/4]$. Then $\text{SFix } T = \{0\}$ see also [10].

Acknowledgments

The authors are grateful to the referee for the professional comments that improved the final version of this paper. Also, the second author was supported by the National University Research Council of the Ministry of Education and Research of Romania, Grant CNCSIS 187.

References

- [1] R. P. Agarwal, J. Dshalalow, and D. O'Regan, "Fixed point and homotopy results for generalized contractive maps of Reich type," *Applicable Analysis*, vol. 82, no. 4, pp. 329–350, 2003.
- [2] A. Chiş and R. Precup, "Continuation theory for general contractions in gauge spaces," *Fixed Point Theory and Applications*, vol. 2004, no. 3, pp. 173–185, 2004.
- [3] L. B. Ćirić, "Generalized contractions and fixed-point theorems," *Publications de l'Institut Mathématique. Nouvelle Série*, vol. 12(26), pp. 19–26, 1971.
- [4] H. Covitz and S. B. Nadler Jr., "Multi-valued contraction mappings in generalized metric spaces," *Israel Journal of Mathematics*, vol. 8, pp. 5–11, 1970.
- [5] Y. Feng and S. Liu, "Fixed point theorems for multi-valued contractive mappings and multi-valued Caristi type mappings," *Journal of Mathematical Analysis and Applications*, vol. 317, no. 1, pp. 103–112, 2006.
- [6] M. Frigon and A. Granas, "Résultats du type de Leray-Schauder pour des contractions multivoques," *Topological Methods in Nonlinear Analysis*, vol. 4, no. 1, pp. 197–208, 1994.
- [7] S. B. Nadler Jr., "Multi-valued contraction mappings," *Pacific Journal of Mathematics*, vol. 30, pp. 475–488, 1969.
- [8] A. Petruşel, "Generalized multivalued contractions," *Nonlinear Analysis*, vol. 47, no. 1, pp. 649–659, 2001.
- [9] A. Petruşel, "Multivalued weakly Picard operators and applications," *Scientiae Mathematicae Japonicae*, vol. 59, no. 1, pp. 169–202, 2004.
- [10] S. Reich, "Fixed points of contractive functions," *Bollettino della Unione Matematica Italiana. Serie IV*, vol. 5, pp. 26–42, 1972.
- [11] S. Reich, "A fixed point theorem for locally contractive multi-valued functions," *Revue Roumaine de Mathématiques Pures et Appliquées*, vol. 17, pp. 569–572, 1972.
- [12] I. A. Rus, "Fixed point theorems for multi-valued mappings in complete metric spaces," *Mathematica Japonica*, vol. 20, pp. 21–24, 1975, special issue.
- [13] I. A. Rus, *Generalized Contractions and Applications*, Cluj University Press, Cluj-Napoca, Romania, 2001.
- [14] I. A. Rus, A. Petruşel, and G. Petruşel, *Fixed Point Theory: 1950–2000. Romanian Contributions*, House of the Book of Science, Cluj-Napoca, Romania, 2002.
- [15] I. A. Rus, A. Petruşel, and A. Sintămărian, "Data dependence of the fixed point set of some multivalued weakly Picard operators," *Nonlinear Analysis*, vol. 52, no. 8, pp. 1947–1959, 2003.

Cristian Chifu: Department of Business, Faculty of Business, Babeş-Bolyai University Cluj-Napoca, Horea 7, 400174 Cluj-Napoca, Romania
 Email address: cochifu@tbs.ubbcluj.ro

Gabriela Petruşel: Department of Applied Mathematics, Faculty of Mathematics and Computer Science, Babeş-Bolyai University Cluj-Napoca, Kogălniceanu 1, 400084 Cluj-Napoca, Romania
 Email address: gabip@math.ubbcluj.ro