

Research Article

Weak and Strong Convergence Theorems for Nonexpansive Mappings in Banach Spaces

Jing Zhao,¹ Songnian He,¹ and Yongfu Su²

¹ College of Science, Civil Aviation University of China, Tianjin 300300, China

² Department of Mathematics, Tianjin Polytechnic University, Tianjin 300160, China

Correspondence should be addressed to Jing Zhao, zhaojing200103@163.com

Received 25 August 2007; Accepted 16 December 2007

Recommended by Tomonari Suzuki

The purpose of this paper is to introduce two implicit iteration schemes for approximating fixed points of nonexpansive mapping T and a finite family of nonexpansive mappings $\{T_i\}_{i=1}^N$, respectively, in Banach spaces and to prove weak and strong convergence theorems. The results presented in this paper improve and extend the corresponding ones of H.-K. Xu and R. Ori, 2001, Z. Opial, 1967, and others.

Copyright © 2008 Jing Zhao et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction and preliminaries

Let E be a real Banach space, K a nonempty closed convex subset of E , and $T : K \rightarrow K$ a mapping. We use $F(T)$ to denote the set of fixed points of T , that is, $F(T) = \{x \in K : Tx = x\}$. T is called nonexpansive if $\|Tx - Ty\| \leq \|x - y\|$ for all $x, y \in K$. In this paper, \rightharpoonup and \rightarrow denote weak and strong convergence, respectively. $\overline{\text{co}}(A)$ denotes the closed convex hull of A , where A is a subset of E .

In 2001, Xu and Ori [1] introduced the following implicit iteration scheme for common fixed points of a finite family of nonexpansive mappings $\{T_i\}_{i=1}^N$ in Hilbert spaces:

$$x_n = \alpha_n x_{n-1} + (1 - \alpha_n) T_n x_n, \quad n \geq 1, \quad (1.1)$$

where $T_n = T_{n \bmod N}$, and they proved weak convergence theorem.

In this paper, we introduce a new implicit iteration scheme:

$$x_n = \alpha_n x_{n-1} + \beta_n T x_{n-1} + \gamma_n T x_n, \quad n \geq 1, \quad (1.2)$$

for fixed points of nonexpansive mapping T in Banach space and also prove weak and strong convergence theorems. Moreover, we introduce an implicit iteration scheme:

$$x_n = \alpha_n x_{n-1} + \beta_n T_n x_{n-1} + \gamma_n T_n x_n, \quad n \geq 1, \quad (1.3)$$

where $T_n = T_{n \bmod N}$, for common fixed points of a finite family of nonexpansive mappings $\{T_i\}_{i=1}^N$ in Banach spaces and also prove weak and strong convergence theorems.

Observe that if K is a nonempty closed convex subset of a real Banach space E and $T : K \rightarrow K$ is a nonexpansive mapping, then for every $u \in K$, $\alpha, \beta, \gamma \in [0, 1]$, and positive integer n , the operator $S = S_{(\alpha, \beta, \gamma, n)} : K \rightarrow K$ defined by

$$Sx = \alpha u + \beta T u + \gamma T x \quad (1.4)$$

satisfies

$$\|Sx - Sy\| = \|\gamma T x - \gamma T y\| \leq \gamma \|x - y\| \quad (1.5)$$

for all $x, y \in K$. Thus, if $\gamma < 1$ then S is a contractive mapping. Then S has a unique fixed point $x^* \in K$. This implies that, if $\gamma_n < 1$, the implicit iteration scheme (1.2) and (1.3) can be employed for the approximation of fixed points of nonexpansive mapping and common fixed points of a finite family of nonexpansive mappings, respectively.

Now, we give some definitions and lemmas for our main results.

A Banach space E is said to satisfy *Opial's condition* if, for any $\{x_n\} \subset E$ with $x_n \rightarrow x \in E$, the following inequality holds:

$$\limsup_{n \rightarrow \infty} \|x_n - x\| < \limsup_{n \rightarrow \infty} \|x_n - y\|, \quad \forall y \in E, \quad x \neq y. \quad (1.6)$$

Let D be a closed subset of a real Banach space E and let $T : D \rightarrow D$ be a mapping.

T is said to be *demiclosed* at zero if $Tx_0 = 0$ whenever $\{x_n\} \subset D$, $x_n \rightarrow x_0$ and $Tx_n \rightarrow 0$.

T is said to be *semicompact* if, for any bounded sequence $\{x_n\} \subset D$ with $\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0$, there exists a subsequence $\{x_{n_k}\} \subset \{x_n\}$ such that $\{x_{n_k}\}$ converges strongly to $x^* \in D$.

Lemma 1.1 (see [2, 3]). *Let E be a uniformly convex Banach space, let K be a nonempty closed convex subset of E , and let $T : K \rightarrow K$ be a nonexpansive mapping. Then $I - T$ is demiclosed at zero.*

Lemma 1.2 (see [4]). *Let E be a uniformly convex Banach space and let a, b be two constants with $0 < a < b < 1$. Suppose that $\{t_n\} \subset [a, b]$ is a real sequence and $\{x_n\}, \{y_n\}$ are two sequences in E . Then the conditions*

$$\lim_{n \rightarrow \infty} \|t_n x_n + (1 - t_n) y_n\| = d, \quad \limsup_{n \rightarrow \infty} \|x_n\| \leq d, \quad \limsup_{n \rightarrow \infty} \|y_n\| \leq d \quad (1.7)$$

imply that $\lim_{n \rightarrow \infty} \|x_n - y_n\| = 0$, where $d \geq 0$ is a constant.

2. Main results

Theorem 2.1. *Let E be a real uniformly convex Banach space which satisfies Opial's condition, let K be a nonempty closed convex subset of E , and let $T : K \rightarrow K$ be a nonexpansive mapping with nonempty fixed points set F . Let $\{\alpha_n\}, \{\beta_n\}, \{\gamma_n\}$ be three real sequences in $[0, 1]$ satisfying $\alpha_n + \beta_n + \gamma_n = 1$ and $0 < a \leq \gamma_n \leq b < 1$, where a, b are some constants. Then implicit iteration process $\{x_n\}$ defined by (1.2) converges weakly to a fixed point of T .*

Proof. Firstly, the condition of Theorem 2.1 implies $\gamma_n < 1$, so that (1.2) can be employed for the approximation of fixed point of nonexpansive mapping.

For any given $p \in F$, we have

$$\begin{aligned}
\|x_n - p\| &= \|\alpha_n x_{n-1} + \beta_n T x_{n-1} + \gamma_n T x_n - p\| \\
&= \|\alpha_n (x_{n-1} - p) + \beta_n (T x_{n-1} - p) + \gamma_n (T x_n - p)\| \\
&\leq \alpha_n \|x_{n-1} - p\| + \beta_n \|T x_{n-1} - p\| + \gamma_n \|T x_n - p\| \\
&\leq (\alpha_n + \beta_n) \|x_{n-1} - p\| + \gamma_n \|x_n - p\|
\end{aligned} \tag{2.1}$$

which leads to

$$(1 - \gamma_n) \|x_n - p\| \leq (\alpha_n + \beta_n) \|x_{n-1} - p\| = (1 - \gamma_n) \|x_{n-1} - p\|. \tag{2.2}$$

It follows from the condition $\gamma_n \leq b < 1$ that

$$\|x_n - p\| \leq \|x_{n-1} - p\|. \tag{2.3}$$

Thus $\lim_{n \rightarrow \infty} \|x_n - p\|$ exists, and so let

$$\lim_{n \rightarrow \infty} \|x_n - p\| = d. \tag{2.4}$$

Hence $\{x_n\}$ is a bounded sequence. Moreover, $\overline{\text{co}}(\{x_n\})$ is a bounded closed convex subset of K . We have

$$\begin{aligned}
\lim_{n \rightarrow \infty} \|x_n - p\| &= \lim_{n \rightarrow \infty} \|\alpha_n (x_{n-1} - p) + \beta_n (T x_{n-1} - p) + \gamma_n (T x_n - p)\| \\
&= \lim_{n \rightarrow \infty} \left\| (1 - \gamma_n) \left[\frac{\alpha_n}{1 - \gamma_n} (x_{n-1} - p) + \frac{\beta_n}{1 - \gamma_n} (T x_{n-1} - p) \right] + \gamma_n (T x_n - p) \right\| = d, \\
\limsup_{n \rightarrow \infty} \|T x_n - p\| &\leq \limsup_{n \rightarrow \infty} \|x_n - p\| = d.
\end{aligned} \tag{2.5}$$

Again, it follows from the condition $\alpha_n + \beta_n + \gamma_n = 1$ that

$$\begin{aligned}
&\limsup_{n \rightarrow \infty} \left\| \frac{\alpha_n}{1 - \gamma_n} (x_{n-1} - p) + \frac{\beta_n}{1 - \gamma_n} (T x_{n-1} - p) \right\| \\
&\leq \limsup_{n \rightarrow \infty} \left(\frac{\alpha_n}{1 - \gamma_n} \|x_{n-1} - p\| + \frac{\beta_n}{1 - \gamma_n} \|T x_{n-1} - p\| \right) \\
&\leq \limsup_{n \rightarrow \infty} \left(\frac{\alpha_n + \beta_n}{1 - \gamma_n} \|x_{n-1} - p\| \right) = d.
\end{aligned} \tag{2.6}$$

By Lemma 1.2, the condition $0 < a \leq \gamma_n \leq b < 1$, and (2.5)–(2.6), we get

$$\lim_{n \rightarrow \infty} \left\| \frac{\alpha_n}{1 - \gamma_n} (x_{n-1} - p) + \frac{\beta_n}{1 - \gamma_n} (T x_{n-1} - p) - (T x_n - p) \right\| = 0. \tag{2.7}$$

This means that

$$\lim_{n \rightarrow \infty} \left\| \frac{\alpha_n}{1 - \gamma_n} x_{n-1} + \frac{\beta_n}{1 - \gamma_n} T x_{n-1} - T x_n \right\| = \lim_{n \rightarrow \infty} \left(\frac{1}{1 - \gamma_n} \right) \|\alpha_n x_{n-1} + \beta_n T x_{n-1} - (1 - \gamma_n) T x_n\| = 0. \quad (2.8)$$

Since $0 < a \leq \gamma_n \leq b < 1$, we have $1/(1 - a) \leq 1/(1 - \gamma_n) \leq 1/(1 - b)$. Hence,

$$\lim_{n \rightarrow \infty} \|\alpha_n x_{n-1} + \beta_n T x_{n-1} - (1 - \gamma_n) T x_n\| = 0. \quad (2.9)$$

Because

$$\begin{aligned} \lim_{n \rightarrow \infty} \|\alpha_n x_{n-1} + \beta_n T x_{n-1} - (1 - \gamma_n) T x_n\| &= \lim_{n \rightarrow \infty} \|x_n - \gamma_n T x_n - (1 - \gamma_n) T x_n\| \\ &= \lim_{n \rightarrow \infty} \|x_n - T x_n\|, \end{aligned} \quad (2.10)$$

by (2.9), we get

$$\lim_{n \rightarrow \infty} \|x_n - T x_n\| = 0. \quad (2.11)$$

Since E is uniformly convex, every bounded closed convex subset of E is weakly compact, so that there exists a subsequence $\{x_{n_k}\}$ of sequence $\{x_n\} \subseteq \overline{\text{co}}(\{x_n\})$ such that $x_{n_k} \rightharpoonup q \in K$. Therefore, it follows from (2.11) that

$$\lim_{k \rightarrow \infty} \|T x_{n_k} - x_{n_k}\| = 0. \quad (2.12)$$

By Lemma 1.1, we know that $I - T$ is demiclosed at zero; it is easy to see that $q \in F$.

Now, we show that $x_n \rightharpoonup q$. In fact, this is not true; then there must exist a subsequence $\{x_{n_i}\} \subset \{x_n\}$ such that $x_{n_i} \rightharpoonup q_1 \in K$, $q_1 \neq q$. Then, by the same method given above, we can also prove that $q_1 \in F$.

Because, for any $p \in F$, the limit $\lim_{n \rightarrow \infty} \|x_n - p\|$ exists. Then we can let

$$\lim_{n \rightarrow \infty} \|x_n - q\| = d_1, \quad \lim_{n \rightarrow \infty} \|x_n - q_1\| = d_2. \quad (2.13)$$

Since E satisfies Opial's condition, we have

$$\begin{aligned} d_1 &= \limsup_{k \rightarrow \infty} \|x_{n_k} - q\| < \limsup_{k \rightarrow \infty} \|x_{n_k} - q_1\| = d_2, \\ d_2 &= \limsup_{i \rightarrow \infty} \|x_{n_i} - q_1\| < \limsup_{i \rightarrow \infty} \|x_{n_i} - q\| = d_1. \end{aligned} \quad (2.14)$$

This is a contradiction and hence $q = q_1$. This implies that $\{x_n\}$ converges weakly to a fixed point q of T . This completes the proof. \square

From the proof of Theorem 2.1, we give the following strong convergence theorem.

Theorem 2.2. *Let E be a real uniformly convex Banach space, let K be a nonempty closed convex subset of E , let $T : K \rightarrow K$ be a nonexpansive mapping with nonempty fixed points set F , and let T be semicompact. Let $\{\alpha_n\}$, $\{\beta_n\}$, $\{\gamma_n\}$ be three real sequences in $[0, 1]$ satisfying $\alpha_n + \beta_n + \gamma_n = 1$ and $0 < a \leq \gamma_n \leq b < 1$, where a, b are some constants. Then implicit iteration process $\{x_n\}$ defined by (1.2) converges strongly to a fixed point of T .*

Proof. From the proof of Theorem 2.1, we know that there exists subsequence $\{x_{n_k}\} \subset \{x_n\}$ such that $x_{n_k} \rightharpoonup q \in K$ and satisfies (2.11). By the semicompactness of T , there exists a subsequence of $\{x_{n_k}\}$ (we still denote it by $\{x_{n_k}\}$) such that $\lim_{n \rightarrow \infty} \|x_{n_k} - q\| = 0$. Because the limit $\lim_{n \rightarrow \infty} \|x_n - q\|$ exists, thus we get $\lim_{n \rightarrow \infty} \|x_n - q\| = 0$. This completes the proof. \square

Next, we study weak and strong convergence theorems for common fixed points of a finite family of nonexpansive mappings $\{T_i\}_{i=1}^N$ in Banach spaces.

Theorem 2.3. *Let E be a real uniformly convex Banach space which satisfies Opial's condition, let K be a nonempty closed convex subset of E , and let $\{T_i\}_{i=1}^N : K \rightarrow K$ be N nonexpansive mappings with nonempty common fixed points set F . Let $\{\alpha_n\}, \{\beta_n\}, \{\gamma_n\}$ be three real sequences in $[0, 1]$ satisfying $\alpha_n + \beta_n + \gamma_n = 1$, $0 < a \leq \gamma_n \leq b < 1$, and $\alpha_n - \beta_n > c > 0$, where a, b, c are some constants. Then implicit iteration process $\{x_n\}$ defined by (1.3) converges weakly to a common fixed point of $\{T_i\}_{i=1}^N$.*

Proof. Substituting T_i ($1 \leq i \leq N$) to T in the proof of Theorem 2.1, we know that for all i ($1 \leq i \leq N$),

$$\lim_{n \rightarrow \infty} \|x_n - T_n x_n\| = 0. \quad (2.15)$$

Now we show that, for any $l = 1, 2, \dots, N$,

$$\lim_{n \rightarrow \infty} \|x_n - T_l x_n\| = 0. \quad (2.16)$$

In fact,

$$\begin{aligned} \|x_n - x_{n-1}\| &= \|\beta_n T_n x_{n-1} + \gamma_n T_n x_n - (\beta_n + \gamma_n)x_{n-1}\| \\ &= \|\beta_n T_n x_{n-1} - \beta_n x_n + \gamma_n T_n x_n - \gamma_n x_n + (\beta_n + \gamma_n)(x_n - x_{n-1})\| \\ &\leq \beta_n \|T_n x_{n-1} - x_n\| + \gamma_n \|T_n x_n - x_n\| + (\beta_n + \gamma_n) \|x_n - x_{n-1}\| \\ &\leq \beta_n \|T_n x_{n-1} - T_n x_n\| + \beta_n \|T_n x_n - x_n\| + \gamma_n \|T_n x_n - x_n\| + (\beta_n + \gamma_n) \|x_n - x_{n-1}\| \\ &\leq (\beta_n + \gamma_n) \|T_n x_n - x_n\| + (2\beta_n + \gamma_n) \|x_n - x_{n-1}\| \\ &= (\beta_n + \gamma_n) \|T_n x_n - x_n\| + (\beta_n + 1 - \alpha_n) \|x_n - x_{n-1}\|. \end{aligned} \quad (2.17)$$

By removing the second term on the right of the above inequality to the left, we get

$$(\alpha_n - \beta_n) \|x_n - x_{n-1}\| \leq (\beta_n + \gamma_n) \|T_n x_n - x_n\|. \quad (2.18)$$

It follows from the condition $\alpha_n - \beta_n > c > 0$ and (2.15) that

$$\lim_{n \rightarrow \infty} \|x_n - x_{n-1}\| = 0. \quad (2.19)$$

So, for any $i = 1, 2, \dots, N$,

$$\lim_{n \rightarrow \infty} \|x_n - x_{n+i}\| = 0. \quad (2.20)$$

Since, for any $i = 1, 2, 3, \dots, N$,

$$\begin{aligned} \|x_n - T_{n+i}x_n\| &\leq \|x_n - x_{n+i}\| + \|x_{n+i} - T_{n+i}x_{n+i}\| + \|T_{n+i}x_{n+i} - T_{n+i}x_n\| \\ &\leq 2\|x_n - x_{n+i}\| + \|x_{n+i} - T_{n+i}x_{n+i}\|, \end{aligned} \quad (2.21)$$

it follows from (2.15) and (2.20) that

$$\lim_{n \rightarrow \infty} \|T_{n+i}x_n - x_n\| = 0, \quad i = 1, 2, 3, \dots, N. \quad (2.22)$$

Because $T_n = T_{n \bmod N}$, it is easy to see, for any $l = 1, 2, 3, \dots, N$, that

$$\lim_{n \rightarrow \infty} \|T_l x_n - x_n\| = 0. \quad (2.23)$$

Since E is uniformly convex, so there exists a subsequence $\{x_{n_k}\}$ of bounded sequence $\{x_n\}$ such that $x_{n_k} \rightharpoonup q \in K$. Therefore, it follows from (2.23) that

$$\lim_{k \rightarrow \infty} \|T_l x_{n_k} - x_{n_k}\| = 0, \quad \forall l = 1, 2, 3, \dots, N. \quad (2.24)$$

By Lemma 1.1, we know that $I - T_l$ is demiclosed, it is easy to see that $q \in F(T_l)$, so that $q \in F = \bigcap_{l=1}^N F(T_l)$. Because E satisfies Opial's condition, we can prove that $\{x_n\}$ converges weakly to a common fixed point q of $\{T_l\}_{l=1}^N$ by the same method given in the proof of Theorem 2.1. \square

Remark 2.4. If $N = 1$, implicit iteration scheme (1.3) becomes (1.2), so from Theorem 2.1, we know that assumption $\alpha_n - \beta_n > c > 0$ in Theorem 2.3 can be removed.

Theorem 2.5. *Let E be a real uniformly convex Banach space, let K be a nonempty closed convex subset of E , let $\{T_i\}_{i=1}^N : K \rightarrow K$ be N nonexpansive mappings with nonempty common fixed points set F , and there exists an $l \in \{1, 2, \dots, N\}$ such that T_l is semicompact. Let $\{\alpha_n\}, \{\beta_n\}, \{\gamma_n\}$ be three real sequences in $[0, 1]$ satisfying $\alpha_n + \beta_n + \gamma_n = 1$, $0 < a \leq \gamma_n \leq b < 1$, and $\alpha_n - \beta_n > c > 0$, where a, b, c are some constants. Then implicit iteration process $\{x_n\}$ defined by (1.3) converges strongly to a common fixed point of $\{T_i\}_{i=1}^N$.*

Proof. From the proof of Theorem 2.3, we know that there exists subsequence $\{x_{n_k}\} \subset \{x_n\}$ such that $\{x_{n_k}\}$ converges weakly to some $q \in K$ and satisfies (2.23). By the semicompactness of T_l , there exists a subsequence of $\{x_{n_k}\}$ (we still denote it by $\{x_{n_k}\}$) such that $\lim_{n \rightarrow \infty} \|x_{n_k} - q\| = 0$. Because the limit $\lim_{n \rightarrow \infty} \|x_n - q\|$ exists, thus we get $\lim_{n \rightarrow \infty} \|x_n - q\| = 0$. This completes the proof. \square

Acknowledgment

This research is supported by Tianjin Natural Science Foundation in China Grant (no. 06YFJMJC12500).

References

- [1] H.-K. Xu and R. G. Ori, "An implicit iteration process for nonexpansive mappings," *Numerical Functional Analysis and Optimization*, vol. 22, no. 5-6, pp. 767–773, 2001.
- [2] H. H. Bauschke, "The approximation of fixed points of compositions of nonexpansive mappings in Hilbert space," *Journal of Mathematical Analysis and Applications*, vol. 202, no. 1, pp. 150–159, 1996.
- [3] J. Górnicki, "Weak convergence theorems for asymptotically nonexpansive mappings in uniformly convex Banach spaces," *Commentationes Mathematicae Universitatis Carolinae*, vol. 30, no. 2, pp. 249–252, 1989.
- [4] J. Schu, "Weak and strong convergence to fixed points of asymptotically nonexpansive mappings," *Bulletin of the Australian Mathematical Society*, vol. 43, no. 1, pp. 153–159, 1991.