

RESEARCH

Open Access

Tripled fixed point theorem in fuzzy metric spaces and applications

A Roldán¹, J Martínez-Moreno^{2*} and C Roldán³

*Correspondence:

jmmoreno@ujaen.es

²Department of Mathematics,
University of Jaén, Jaén, Spain
Full list of author information is
available at the end of the article

Abstract

In this paper we prove an existence and uniqueness theorem for contractive type mappings in fuzzy metric spaces. In order to do that, we consider a slight modification of the concept of a tripled fixed point introduced by Berinde *et al.* (Nonlinear Anal. TMA 74:4889-4897, 2011) for nonlinear mappings. Additionally, we obtain some fixed point theorems for metric spaces. These results generalize, extend and unify several classical and very recent related results in literature. For instance, we obtain an extension of Theorem 4.1 in (Zhu and Xiao in Nonlinear Anal. TMA 74:5475-5479, 2011) and a version in non-partially ordered sets of Theorem 2.2 in (Bhaskar and Lakshmikantham in Nonlinear Anal. TMA 65:1379-1393, 2006). As application, we solve a kind of Lipschitzian systems in three variables and an integral system. Finally, examples to support our results are also given.

Introduction

In a recent paper, Bhaskar and Lakshmikantham [1] introduced the concepts of *coupled fixed point* and *mixed monotone property* for contractive operators of the form $F : X \times X \rightarrow X$, where X is a partially ordered metric space, and then established some interesting coupled fixed point theorems. They also illustrated these important results by proving the existence and uniqueness of the solution for a periodic boundary value problem. Later, Lakshmikantham and Ćirić [2] proved coupled coincidence and coupled common fixed point results for nonlinear mappings satisfying certain contractive conditions in partially ordered complete metric spaces. After that many results appeared on coupled fixed point theory (see, *e.g.*, [2–8]).

Fixed point theorems have been studied in many contexts, one of which is the fuzzy setting. The concept of *fuzzy sets* was initially introduced by Zadeh [9] in 1965. To use this concept in topology and analysis, many authors have extensively developed the theory of fuzzy sets and its applications. One of the most interesting research topics in fuzzy topology is to find an appropriate definition of *fuzzy metric space* for its possible applications in several areas. It is well known that a fuzzy metric space is an important generalization of the metric space. Many authors have considered this problem and have introduced it in different ways. For instance, George and Veeramani [10] modified the concept of a fuzzy metric space introduced by Kramosil and Michalek [11] and defined the Hausdorff topology of a fuzzy metric space. There exists considerable literature about fixed point properties for mappings defined on fuzzy metric spaces, which have been studied by many authors (see [10, 12–16]). Zhu and Xiao [7] and Hu [6] gave a coupled fixed point theorem for contractions in fuzzy metric spaces, and Fang [3] proved some common fixed

point theorems under ϕ -contractions for compatible and weakly compatible mappings on Menger probabilistic metric spaces. Moreover, Elagan and Segi Rahmat [17] studied the existence of a fixed point in locally convex topology generated by fuzzy n -normed spaces.

Very recently, the concept of *tripled fixed point* has been introduced by Berinde and Borcut [18]. In their manuscript, some new tripled point theorems are obtained using the mixed g -monotone mapping. Their results generalize and extend the Bhaskar and Lakshmikantham's research for nonlinear mappings. Moreover, these results could be used to study the existence of solutions of a periodic boundary value problem involving $y'' = f(t, y, y')$. A multidimensional notion of a coincidence point between mappings and some existence and uniqueness fixed points theorems for nonlinear mappings defined on partially ordered metric spaces are studied in [19].

In this paper, our main aim is to obtain an existence and uniqueness theorem for contractive type mappings in the framework of fuzzy metric spaces. In order to do that, we consider a slight modification of the concept of a tripled fixed point introduced by Berinde and Borcut for nonlinear mappings. The power of this result is two-fold. Firstly, we can particularize it to complete metric spaces, obtaining a Berinde-Borcut type result (in non-fuzzy setting). Moreover, our result, in a unified manner, covers also coupled fixed (see Zhu and Xiao [7]) and fixed point theorems. Finally, examples to support our results are also given.

Preliminaries

Henceforth, X will denote a non-empty set and $X^3 = X \times X \times X$. Subscripts will be used to indicate the arguments of a function. For instance, $F(x, y, z)$ will be denoted by F_{xyz} and $M(x, y, t)$ will be denoted by $M_{xy}(t)$. Furthermore, for brevity, $g(x)$ will be denoted by gx .

A *metric on X* is a mapping $d : X \times X \rightarrow \mathbb{R}$ satisfying, for all $x, y, z \in X$,

- (i) $d_{xy} = 0$ if and only if $x = y$;
- (ii) $d_{xy} \leq d_{zx} + d_{zy}$.

From these properties, we can easily deduce that $d_{xy} \geq 0$ and $d_{yx} = d_{xy}$ for all $x, y \in X$. The last requirement is called the *triangle inequality*. If d is a metric on X , we say that (X, d) is a *metric space* (briefly, a *MS*).

Let (X, d) be a MS. A mapping $f : X \rightarrow X$ is said to be *Lipschitzian* if there exists $k \geq 0$ such that $d(f_x, f_y) \leq kd_{xy}$ for all $x, y \in X$. The smallest k (denoted by k_f) for which this inequality holds is said to be the *Lipschitz constant for f* . A Lipschitzian mapping $f : X \rightarrow X$ is a *contraction* if $k_f < 1$.

Theorem 1 (Banach's contraction principle) *Every contraction from a complete metric space into itself has a unique fixed point.*

If $X = \mathbb{R}$ provided with the Euclidean metric, examples of Lipschitzian mappings $f_i : \mathbb{R} \rightarrow \mathbb{R}$ are $f_1(x) = K, f_2(x) = \alpha x, f_3(x) = \sin x, f_4(x) = \cos x, f_5(x) = \arctan x$ and $f_6(x) = 1/(1 + x^2)$.

Definition 2 A *triangular norm* (also called a *t-norm*) is a map $* : [0, 1]^2 \rightarrow [0, 1]$ that is associative, commutative, nondecreasing in both arguments and has 1 as identity. For each $a \in [0, 1]$, the sequence $\{ *^n a \}_{n=1}^\infty$ is defined inductively by $*^1 a = a$ and $*^n a = (*^{n-1} a) * a$. A *t-norm* $*$ is said to be of *H-type* (see [20]) if the sequence $\{ *^n a \}_{n=1}^\infty$ is equicontinuous at

$a = 1$, i.e., for all $\varepsilon \in (0, 1)$, there exists $\eta \in (0, 1)$ such that if $a \in (1 - \eta, 1]$, then $*^m a > 1 - \varepsilon$ for all $m \in \mathbb{N}$.

The most important and well-known continuous t -norm of H -type is $* = \min$, that verifies $\min(a, b) \geq ab$ for all $a, b \in [0, 1]$. The following result presents a wide range of t -norms of H -type.

Lemma 3 Let $\delta \in (0, 1]$ be a real number and let $*$ be a t -norm. Define $*_\delta$ as $x *_\delta y = x * y$, if $\max(x, y) \leq 1 - \delta$, and $x *_\delta y = \min(x, y)$, if $\max(x, y) > 1 - \delta$. Then $*_\delta$ is a t -norm of H -type.

Definition 4 [11] A triple $(X, M, *)$ is called a *fuzzy metric space* (in the sense of Kramosil and Michalek; briefly, a FMS) if X is an arbitrary non-empty set, $*$ is a continuous t -norm and $M : X \times X \times [0, \infty) \rightarrow [0, 1]$ is a fuzzy set satisfying the following conditions, for each $x, y, z \in X$, and $t, s > 0$:

- (i) $M_{xy}(0) = 0$;
- (ii) $M_{xy}(t) = 1$ if and only if $x = y$;
- (iii) $M_{xy}(t) = M_{yx}(t)$;
- (iv) $M_{xy}(\cdot) : [0, \infty) \rightarrow [0, 1]$ is left continuous;
- (v) $M_{xy}(t) * M_{yz}(s) \leq M_{xz}(t + s)$.

In this case, we also say that (X, M) is a FMS under $*$. In the sequel, we will only consider FMS verifying:

- (vi) $\lim_{t \rightarrow \infty} M_{xy}(t) = 1$ for all $x, y \in X$.

Lemma 5 $M_{xy}(\cdot)$ is a non-decreasing function on $[0, \infty)$.

Definition 6 Let (X, M) be a FMS under some t -norm. A sequence $\{x_n\} \subset X$ is *Cauchy* if, for any $\varepsilon > 0$ and $t > 0$, there exists $n_0 \in \mathbb{N}$ such that $M_{x_n, x_m}(t) > 1 - \varepsilon$ for all $n, m \geq n_0$. A sequence $\{x_n\} \subset X$ is *convergent to* $x \in X$, denoted by $\lim_{n \rightarrow \infty} x_n = x$ if, for any $\varepsilon > 0$ and $t > 0$, there exists $n_0 \in \mathbb{N}$ such that $M_{x_n, x}(t) > 1 - \varepsilon$ for all $n \geq n_0$. A FMS in which every Cauchy sequence is convergent is called *complete*.

Given any t -norm $*$, it is easy to prove that $* \leq \min$. Therefore, if (X, M) is a FMS under \min , then (X, M) is a FMS under any (continuous or not) t -norm. This is the case in the following examples (in which, obviously, we only define $M_{xy}(t)$ for $t > 0$ and $x \neq y$).

Example 7 From a metric space (X, d) , we can consider a FMS in different ways. For $t > 0$ and $x \neq y$, define:

$$\bullet \quad M_{xy}^d(t) = \frac{t}{t + d_{xy}} \quad \bullet \quad M_{xy}^e(t) = e^{-\frac{d_{xy}}{t}} \quad \bullet \quad M_{x,y}^c(t) = \begin{cases} 0, & \text{if } t \leq d_{xy}, \\ 1, & \text{if } t > d_{xy}. \end{cases}$$

It is well known that (X, M^d) is a FMS under the product $* = \cdot$, called the *standard FMS on* (X, d) , since it is the standard way of viewing the metric space (X, d) as a FMS. However, it is also true (though lesser-known) that (X, M^d) , (X, M^e) and (X, M^c) are FMSs under \min .

Furthermore, (X, d) is a complete metric space if and only if (X, M^d) (or (X, M^c) or (X, M^e)) is a complete FMS. For instance, this is the case of any non-empty and closed subset (or subinterval) of \mathbb{R} provided with its Euclidean metric.

Definition 8 A function $g : X \rightarrow X$ on a FMS is said to be *continuous at a point* $x_0 \in X$ if, for any sequence $\{x_n\}$ in X converging to x_0 , the sequence $\{gx_n\}$ converges to gx_0 . If g is continuous at each $x \in X$, then g is said to be *continuous on* X . As usual, if $x_0 \in X$, we will denote $g^{-1}(x_0) = \{x \in X : gx = x_0\}$.

Remark 9 If $x \in [0, 1]$ and $a, b \in (0, \infty)$, then $a \leq b$ implies that $x^a \geq x^b$. We will use this fact in the following way: $0 < a \leq b \leq 1$ implies that $M_{xy}(t)^a \geq M_{xy}(t)^b \geq M_{xy}(t)$.

The main result

Definition 10 Let $F : X^3 \rightarrow X$ and $g : X \rightarrow X$ be two mappings.

- We say that F and g are *commuting* if $gF_{xyz} = F_{gxygz}$ for all $x, y, z \in X$.
- A point $(x, y, z) \in X^3$ is called a *tripled coincidence point of the mappings* F and g if $F_{xyz} = gx, F_{yzx} = gy$ and $F_{zxy} = gz$.

Theorem 11 Let $*$ be a t -norm of H -type such that $s * t \geq st$ for all $s, t \in [0, 1]$. Let $k \in (0, 1)$ and $a, b, c \in [0, 1]$ be real numbers such that $a + b + c \leq 1$, let $(X, M, *)$ be a complete FMS and let $F : X^3 \rightarrow X$ and $g : X \rightarrow X$ be two mappings such that $F(X^3) \subseteq g(X)$ and g is continuous and commuting with F . Suppose that for all $x, y, z, u, v, w \in X$ and all $t > 0$,

$$M_{F_{xyz}F_{uvw}}(kt) \geq M_{g_xgu}(t)^a * M_{g_ygv}(t)^b * M_{g_zgw}(t)^c. \tag{1}$$

Then there exists a unique $x \in X$ such that $x = gx = F_{xxx}$. In particular, F and g have, at least, one tripled coincidence point. Furthermore, (x, x, x) is the unique tripled coincidence point of F and g if we assume that $g^{-1}(x_0) = \{x_0\}$ only in the case that $F \equiv x_0$ is constant on X^3 .

In this result, in order to avoid the indetermination 0^0 , we assume that $M_{g_xgu}(t)^0 = 1$ for all $t > 0$ and all $x, y \in X$.

Proof Suppose that F is constant in X^3 , i.e., there exists $x_0 \in X$ such that $F_{xyz} = x_0$ for all $x, y, z \in X$. As F and g are commuting, we deduce that $gx_0 = gF_{xyz} = F_{gxygz} = x_0$. Therefore, $x_0 = gx_0 = F_{x_0x_0x_0}$ and (x_0, x_0, x_0) is a tripled coincidence point of F and g . Now, suppose that $g^{-1}(x_0) = \{x_0\}$ and $(x, y, z) \in X^3$ is another tripled coincidence point of F and g . Then $gx = F_{xyz} = x_0$, so $x \in g^{-1}(x_0) = \{x_0\}$. Similarly, $x = y = z = x_0$ and (x_0, x_0, x_0) is the unique tripled coincidence point of F and g .

Next, suppose that F is not constant in X^3 . In this case, $(a, b, c) \neq (0, 0, 0)$ and the proof is divided into five steps. Throughout this proof, n and p will denote non-negative integers and $t \in [0, \infty)$.

Step 1. Definition of the sequences $\{x_n\}$, $\{y_n\}$ and $\{z_n\}$. Let $x_0, y_0, z_0 \in X$ be three arbitrary points of X . Since $F(X^3) \subseteq g(X)$, we can choose $x_1, y_1, z_1 \in X$ such that $gx_1 = F_{x_0y_0z_0}$, $gy_1 = F_{y_0z_0x_0}$ and $gz_1 = F_{z_0x_0y_0}$. Again, from $F(X^3) \subseteq g(X)$, we can choose $x_2, y_2, z_2 \in X$ such that $gx_2 = F_{x_1y_1z_1}$, $gy_2 = F_{y_1z_1x_1}$ and $gz_2 = F_{z_1x_1y_1}$. Continuing this process, we can construct sequences $\{x_n\}$, $\{y_n\}$ and $\{z_n\}$ such that, for $n \geq 0$, $gx_{n+1} = F_{x_ny_nz_n}$, $gy_{n+1} = F_{y_nz_nx_n}$ and $gz_{n+1} = F_{z_nx_ny_n}$.

Step 2. $\{gx_n\}$, $\{gy_n\}$ and $\{gz_n\}$ are Cauchy sequences. Define, for $n \geq 0$ and all $t \geq 0$, $\delta_n(t) = M_{g_xn g_{x_{n+1}}}(t) * M_{g_yn g_{y_{n+1}}}(t) * M_{g_zn g_{z_{n+1}}}(t)$. Since δ_n is a non-decreasing function and $t - kt \leq$

$t \leq t/k$, we have that

$$\delta_n(t - kt) \leq \delta_n(t) \leq \delta_n(t/k) \quad \text{for all } t > 0 \text{ and } n \geq 0. \tag{2}$$

From inequality (1) we deduce, for all $n \in \mathbb{N}$ and all $t \geq 0$,

$$\begin{aligned} M_{g^{x_n}g^{x_{n+1}}}(t) &= M_{F_{x_{n-1}y_{n-1}z_{n-1}F_{x_ny_nz_n}}}(t) \\ &\geq M_{g^{x_{n-1}g^{x_n}}}\left(\frac{t}{k}\right)^a * M_{g^{y_{n-1}g^{y_n}}}\left(\frac{t}{k}\right)^b * M_{g^{z_{n-1}g^{z_n}}}\left(\frac{t}{k}\right)^c; \end{aligned} \tag{3}$$

$$\begin{aligned} M_{g^{y_n}g^{y_{n+1}}}(t) &= M_{F_{y_{n-1}z_{n-1}x_{n-1}F_{y_nz_nx_n}}}(t) \\ &\geq M_{g^{y_{n-1}g^{y_n}}}\left(\frac{t}{k}\right)^a * M_{g^{z_{n-1}g^{z_n}}}\left(\frac{t}{k}\right)^b * M_{g^{x_{n-1}g^{x_n}}}\left(\frac{t}{k}\right)^c; \end{aligned} \tag{4}$$

$$\begin{aligned} M_{g^{z_n}g^{z_{n+1}}}(t) &= M_{F_{z_{n-1}x_{n-1}y_{n-1}F_{z_nx_ny_n}}}(t) \\ &\geq M_{g^{z_{n-1}g^{z_n}}}\left(\frac{t}{k}\right)^a * M_{g^{x_{n-1}g^{x_n}}}\left(\frac{t}{k}\right)^b * M_{g^{y_{n-1}g^{y_n}}}\left(\frac{t}{k}\right)^c. \end{aligned} \tag{5}$$

According to (3), (4), (5) and Remark 9, we have that

$$\begin{aligned} M_{g^{x_n}g^{x_{n+1}}}(t) &\geq M_{g^{x_{n-1}g^{x_n}}}(t/k)^a * M_{g^{y_{n-1}g^{y_n}}}(t/k)^b * M_{g^{z_{n-1}g^{z_n}}}(t/k)^c \\ &\geq M_{g^{x_{n-1}g^{x_n}}}(t/k) * M_{g^{y_{n-1}g^{y_n}}}(t/k) * M_{g^{z_{n-1}g^{z_n}}}(t/k) = \delta_{n-1}(t/k); \\ M_{g^{y_n}g^{y_{n+1}}}(t) &\geq M_{g^{y_{n-1}g^{y_n}}}(t/k)^a * M_{g^{z_{n-1}g^{z_n}}}(t/k)^b * M_{g^{x_{n-1}g^{x_n}}}(t/k)^c \\ &\geq M_{g^{y_{n-1}g^{y_n}}}(t/k) * M_{g^{z_{n-1}g^{z_n}}}(t/k) * M_{g^{x_{n-1}g^{x_n}}}(t/k) = \delta_{n-1}(t/k); \\ M_{g^{z_n}g^{z_{n+1}}}(t) &\geq M_{g^{z_{n-1}g^{z_n}}}(t/k)^a * M_{g^{x_{n-1}g^{x_n}}}(t/k)^b * M_{g^{y_{n-1}g^{y_n}}}(t/k)^c \\ &\geq M_{g^{z_{n-1}g^{z_n}}}(t/k) * M_{g^{x_{n-1}g^{x_n}}}(t/k) * M_{g^{y_{n-1}g^{y_n}}}(t/k) = \delta_{n-1}(t/k). \end{aligned}$$

This proves that, for all $t > 0$ and all $n \geq 0$,

$$M_{g^{x_n}g^{x_{n+1}}}(t), M_{g^{y_n}g^{y_{n+1}}}(t), M_{g^{z_n}g^{z_{n+1}}}(t) \geq \delta_{n-1}(t/k) \geq \delta_{n-1}(t). \tag{6}$$

Swapping t by $t - kt$, we deduce, for all $t > 0$ and $n \geq 0$, that

$$M_{g^{x_n}g^{x_{n+1}}}(t - kt), M_{g^{y_n}g^{y_{n+1}}}(t - kt), M_{g^{z_n}g^{z_{n+1}}}(t - kt) \geq \delta_{n-1}(t - kt). \tag{7}$$

Taking into account that $*$ is commutative and $* \geq \cdot$, and (3), (4), (5), we observe that

$$\begin{aligned} \delta_n(t) &= M_{g^{x_n}g^{x_{n+1}}}(t) * M_{g^{y_n}g^{y_{n+1}}}(t) * M_{g^{z_n}g^{z_{n+1}}}(t) \\ &\geq \left(M_{g^{x_{n-1}g^{x_n}}}(t/k)^a * M_{g^{y_{n-1}g^{y_n}}}(t/k)^b * M_{g^{z_{n-1}g^{z_n}}}(t/k)^c \right) \\ &\quad * \left(M_{g^{x_{n-1}g^{x_n}}}(t/k)^c * M_{g^{y_{n-1}g^{y_n}}}(t/k)^a * M_{g^{z_{n-1}g^{z_n}}}(t/k)^b \right) \\ &\quad * \left(M_{g^{x_{n-1}g^{x_n}}}(t/k)^b * M_{g^{y_{n-1}g^{y_n}}}(t/k)^c * M_{g^{z_{n-1}g^{z_n}}}(t/k)^a \right) \\ &= \left(M_{g^{x_{n-1}g^{x_n}}}(t/k)^a * M_{g^{x_{n-1}g^{x_n}}}(t/k)^c * M_{g^{x_{n-1}g^{x_n}}}(t/k)^b \right) \\ &\quad * \left(M_{g^{y_{n-1}g^{y_n}}}(t/k)^b * M_{g^{y_{n-1}g^{y_n}}}(t/k)^a * M_{g^{x_{n-1}g^{x_n}}}(t/k)^c \right) \end{aligned}$$

$$\begin{aligned}
 & * (M_{gz_{n-1}gz_n}(t/k)^c * M_{gz_{n-1}gz_n}(t/k)^b * M_{gz_{n-1}gz_n}(t/k)^a) \\
 \geq & (M_{gx_{n-1}gx_n}(t/k)^a \cdot M_{gx_{n-1}gx_n}(t/k)^c \cdot M_{gx_{n-1}gx_n}(t/k)^b) \\
 & * (M_{gy_{n-1}gy_n}(t/k)^b \cdot M_{gy_{n-1}gy_n}(t/k)^a \cdot M_{gx_{n-1}gx_n}(t/k)^c) \\
 & * (M_{gz_{n-1}gz_n}(t/k)^c \cdot M_{gz_{n-1}gz_n}(t/k)^b \cdot M_{gz_{n-1}gz_n}(t/k)^a) \\
 = & M_{gx_{n-1}gx_n}(t/k)^{a+b+c} * M_{gy_{n-1}gy_n}(t/k)^{a+b+c} * M_{gz_{n-1}gz_n}(t/k)^{a+b+c} \\
 \geq & M_{gx_{n-1}gx_n}(t/k) * M_{gy_{n-1}gy_n}(t/k) * M_{gz_{n-1}gz_n}(t/k) = \delta_{n-1}(t/k).
 \end{aligned}$$

If we join this property to (2),

$$\delta_n(t) \geq \delta_{n-1}(t/k) \geq \delta_{n-1}(t) \geq \delta_{n-1}(t - kt) \quad \text{for all } t > 0 \text{ and } n \geq 1. \tag{8}$$

Repeatedly applying the first inequality, we deduce that $\delta_n(t) \geq \delta_{n-1}(t/k) \geq \delta_{n-2}(t/k^2) \geq \dots \geq \delta_0(t/k^n)$ for all $t > 0$ and $n \geq 1$. This means that for all $t > 0$,

$$\lim_{n \rightarrow \infty} \delta_n(t) \geq \lim_{n \rightarrow \infty} \delta_0(t/k^n) = 1 \quad \Rightarrow \quad \lim_{n \rightarrow \infty} \delta_n(t) = 1. \tag{9}$$

Properties (6) and (8) imply that

$$M_{gx_n gx_{n+1}}(t), M_{gy_n gy_{n+1}}(t), M_{gz_n gz_{n+1}}(t) \geq \delta_n(t) \geq \delta_{n-1}(t - kt). \tag{10}$$

Next, we claim that

$$M_{gx_n gx_{n+p}}(t), M_{gy_n gy_{n+p}}(t), M_{gz_n gz_{n+p}}(t) \geq *^p \delta_{n-1}(t - kt) \quad \text{for all } t > 0, n, p \geq 1. \tag{11}$$

We prove it by induction methodology in $p \geq 1$. If $p = 1$, (11) is true for all $n \geq 1$ and all $t > 0$ by (10). Suppose that (11) is true for all $n \geq 1$ and all $t > 0$ for some p , and we are going to prove it for $p + 1$. Applying (1), the induction hypothesis and that $* \geq \cdot$,

$$\begin{aligned}
 M_{gx_{n+1}gx_{n+p+1}}(kt) & = M_{\Gamma_{x_n y_n z_n} F_{x_{n+p} y_{n+p} z_{n+p}}}(kt) \\
 & \geq M_{gx_n gx_{n+p}}(t)^a * M_{gy_n gy_{n+p}}(t)^b * M_{gz_n gz_{n+p}}(t)^c \\
 & \geq (*^p \delta_{n-1}(t - kt))^a * (*^p \delta_{n-1}(t - kt))^b * (*^p \delta_{n-1}(t - kt))^c \\
 & \geq (*^p \delta_{n-1}(t - kt))^a \cdot (*^p \delta_{n-1}(t - kt))^b \cdot (*^p \delta_{n-1}(t - kt))^c \\
 & = (*^p \delta_{n-1}(t - kt))^{a+b+c} \geq *^p \delta_{n-1}(t - kt).
 \end{aligned}$$

Arguing in the same way, we come to $M_{gx_{n+1}gx_{n+1+p}}(kt), M_{gy_{n+1}gy_{n+1+p}}(kt), M_{gz_{n+1}gz_{n+1+p}}(kt) \geq *^p \delta_{n-1}(t - kt)$. Applying the axiom (v) of a FMS, (7) and the induction hypothesis,

$$\begin{aligned}
 M_{gx_n gx_{n+p+1}}(t) & = M_{gx_n gx_{n+p+1}}(t - kt + kt) \\
 & \geq M_{gx_n gx_{n+1}}(t - kt) * M_{gx_{n+1}gx_{n+1+p}}(kt) \\
 & \geq \delta_{n-1}(t - kt) * (*^p \delta_{n-1}(t - kt)) = *^{p+1} \delta_{n-1}(t - kt).
 \end{aligned}$$

The same reasoning is also valid for $M_{g_n g_{n+p+1}}(t)$ and $M_{g_n g_{n+p+1}}(t)$. Therefore, (11) is true. This permits us to show that $\{g x_n\}$ is Cauchy. Suppose that $t > 0$ and $\varepsilon \in (0, 1)$ are given. By the hypothesis, as $*$ is a t -norm of H -type, there exists $0 < \eta < 1$ such that $*^p a > 1 - \varepsilon$ for all $a \in (1 - \eta, 1]$ and for all $p \geq 1$. By (9), $\lim_{n \rightarrow \infty} \delta_n(t) = 1$, so there exists $n_0 \in \mathbb{N}$ such that $\delta_n(t - kt) > 1 - \eta$ for all $n \geq n_0$. Hence (11), we get $M_{g_n g_{n+p}}(t), M_{g_n g_{n+p}}(t), M_{g_n g_{n+p}}(t) > 1 - \varepsilon$ for all $n \geq n_0$ and $p \geq 1$. Therefore, $\{g x_n\}$ is a Cauchy sequence. Similarly, $\{g y_n\}$ and $\{g z_n\}$ are also Cauchy sequences.

Step 3. We claim that g and F have a tripled coincidence point. Since X is complete, there exist $x, y, z \in X$ such that $\lim_{n \rightarrow \infty} g x_n = x, \lim_{n \rightarrow \infty} g y_n = y$ and $\lim_{n \rightarrow \infty} g z_n = z$. As g is continuous, we have that $\lim_{n \rightarrow \infty} g g x_n = g x, \lim_{n \rightarrow \infty} g g y_n = g y$ and $\lim_{n \rightarrow \infty} g g z_n = g z$. The commutativity of F with g implies that $g g x_{n+1} = g F(x_n, y_n, z_n) = F(g x_n, g y_n, g z_n)$. By (1),

$$\begin{aligned} M_{g g x_{n+1} F_{xyz}}(kt) &= M_{F_{g x_n g y_n g z_n} F_{xyz}}(kt) \geq M_{g g x_n g x}(t)^a * M_{g g y_n g y}(t)^b * M_{g g z_n g z}(t)^c \\ &\geq M_{g g x_n g x}(t) * M_{g g y_n g y}(t) * M_{g g z_n g z}(t). \end{aligned}$$

Letting $n \rightarrow \infty$, we deduce that $\lim_{n \rightarrow \infty} g g x_n = F_{xyz}$. Hence, $F_{xyz} = g x$. In a similar way, we can show that $F_{yzx} = g y$ and $F_{zxy} = g z$, so (x, y, z) is a tripled coincidence point of the mappings F and g .

$$F_{xyz} = g x, \quad F_{yzx} = g y \quad \text{and} \quad F_{zxy} = g z. \tag{12}$$

Step 4. We claim that $x = F_{zxy}, y = F_{xyz}$ and $z = F_{yzx}$. We note that by condition (1),

$$M_{g x g y_{n+1}}(kt) = M_{F_{xyz} F_{yzx} F_{zxy}}(kt) \geq M_{g x g y_n}(t)^a * M_{g y g z_n}(t)^b * M_{g z g x_n}(t)^c; \tag{13}$$

$$M_{g y g z_{n+1}}(kt) = M_{F_{yzx} F_{zxy} F_{xyz}}(kt) \geq M_{g y g z_n}(t)^a * M_{g z g x_n}(t)^b * M_{g x g y_n}(t)^c; \tag{14}$$

$$M_{g z g x_{n+1}}(kt) = M_{F_{zxy} F_{xyz} F_{yzx}}(kt) \geq M_{g z g x_n}(t)^a * M_{g x g y_n}(t)^b * M_{g y g z_n}(t)^c. \tag{15}$$

Let $\beta_n(t) = M_{g x g y_n}(t) * M_{g y g z_n}(t) * M_{g z g x_n}(t)$ for all $t > 0$ and $n \geq 0$. By (13), (14) and (15),

$$\begin{aligned} \beta_{n+1}(kt) &= M_{g x g y_{n+1}}(kt) * M_{g y g z_{n+1}}(kt) * M_{g z g x_{n+1}}(kt) \\ &\geq (M_{g x g y_n}(t)^a * M_{g y g z_n}(t)^b * M_{g z g x_n}(t)^c) \\ &\quad * (M_{g y g z_n}(t)^a * M_{g z g x_n}(t)^b * M_{g x g y_n}(t)^c) \\ &\quad * (M_{g z g x_n}(t)^a * M_{g x g y_n}(t)^b * M_{g y g z_n}(t)^c) \\ &= (M_{g x g y_n}(t)^a * M_{g x g y_n}(t)^c * M_{g x g y_n}(t)^b) \\ &\quad * (M_{g y g z_n}(t)^b * M_{g y g z_n}(t)^a * M_{g y g z_n}(t)^c) \\ &\quad * (M_{g z g x_n}(t)^c * M_{g z g x_n}(t)^b * M_{g z g x_n}(t)^a) \\ &\geq (M_{g x g y_n}(t)^a \cdot M_{g x g y_n}(t)^c \cdot M_{g x g y_n}(t)^b) \\ &\quad * (M_{g y g z_n}(t)^b \cdot M_{g y g z_n}(t)^a \cdot M_{g y g z_n}(t)^c) \\ &\quad * (M_{g z g x_n}(t)^c \cdot M_{g z g x_n}(t)^b \cdot M_{g z g x_n}(t)^a) \\ &= M_{g x g y_n}(t)^{a+b+c} * M_{g y g z_n}(t)^{a+b+c} * M_{g z g x_n}(t)^{a+b+c} \\ &\geq M_{g x g y_n}(t) * M_{g y g z_n}(t) * M_{g z g x_n}(t) = \beta_n(t). \end{aligned}$$

This proves that $\beta_{n+1}(kt) \geq \beta_n(t)$ for all $n \geq 0$ and all $t > 0$. Repeating this process,

$$\beta_n(t) \geq \beta_{n-1}(t/k) \geq \beta_{n-2}(t/k^2) \geq \dots \geq \beta_0(t/k^n) \quad \text{for all } t > 0 \text{ and } n \geq 1. \quad (16)$$

Now, by (16), (13), (14) and (15),

$$M_{g_x g_y g_{n+1}}(kt) \geq M_{g_x g_y g_n}(t)^a * M_{g_y g_z g_n}(t)^b * M_{g_z g_x g_n}(t)^c \geq \beta_n(t) \geq \beta_0(t/k^n); \quad (17)$$

$$M_{g_y g_z g_{n+1}}(kt) \geq M_{g_y g_z g_n}(t)^a * M_{g_z g_x g_n}(t)^b * M_{g_x g_y g_n}(t)^c \geq \beta_n(t) \geq \beta_0(t/k^n); \quad (18)$$

$$M_{g_z g_x g_{n+1}}(kt) \geq M_{g_z g_x g_n}(t)^a * M_{g_x g_y g_n}(t)^b * M_{g_y g_z g_n}(t)^c \geq \beta_n(t) \geq \beta_0(t/k^n). \quad (19)$$

Therefore, $M_{g_x g_y g_{n+1}}(kt), M_{g_y g_z g_{n+1}}(kt), M_{g_z g_x g_{n+1}}(kt) \geq \beta_0(t/k^n)$ for all $t > 0$ and $n \geq 1$. Since $\lim_{n \rightarrow \infty} \beta_0(t/k^n) = 1$ for all $t > 0$, we have, taking limit in (17), (18) and (19), that $\lim_{n \rightarrow \infty} g x_n = g z, \lim_{n \rightarrow \infty} g y_n = g x$ and $\lim_{n \rightarrow \infty} g z_n = g y$. This shows, using (12), that

$$F_{xyz} = g x = \lim_{n \rightarrow \infty} g y_n = y, \quad F_{yzx} = g y = \lim_{n \rightarrow \infty} g z_n = z, \quad F_{zxy} = g z = \lim_{n \rightarrow \infty} g x_n = x.$$

Step 5. We will prove that $x = y = z$. Let $\theta(t) = M_{xy}(t) * M_{yz}(t) * M_{zx}(t)$ for all $t > 0$. Then, by condition (1),

$$\begin{aligned} M_{xy}(kt) &= M_{F_{xyz} F_{yzx}}(kt) \geq M_{g_x g_y}(t)^a * M_{g_y g_z}(t)^b * M_{g_z g_x}(t)^c \\ &= M_{yz}(t)^a * M_{zx}(t)^b * M_{xy}(t)^c; \end{aligned} \quad (20)$$

$$\begin{aligned} M_{yz}(kt) &= M_{F_{yzx} F_{zxy}}(kt) \geq M_{g_y g_z}(t)^a * M_{g_z g_x}(t)^b * M_{g_x g_y}(t)^c \\ &= M_{zx}(t)^a * M_{xy}(t)^b * M_{yz}(t)^c; \end{aligned} \quad (21)$$

$$\begin{aligned} M_{zx}(kt) &= M_{F_{zxy} F_{xyx}}(kt) \geq M_{g_z g_x}(t)^a * M_{g_x g_y}(t)^b * M_{g_y g_z}(t)^c \\ &= M_{xy}(t)^a * M_{yz}(t)^b * M_{zx}(t)^c. \end{aligned} \quad (22)$$

If we use these three inequalities at the same time,

$$\begin{aligned} \theta(kt) &= M_{xy}(kt) * M_{yz}(kt) * M_{zx}(kt) \\ &\geq (M_{yz}(t)^a * M_{zx}(t)^b * M_{xy}(t)^c) * (M_{zx}(t)^a * M_{xy}(t)^b * M_{yz}(t)^c) \\ &\quad * (M_{xy}(t)^a * M_{yz}(t)^b * M_{zx}(t)^c) \\ &= (M_{xy}(t)^c * M_{xy}(t)^b * M_{xy}(t)^a) * (M_{yz}(t)^a * M_{yz}(t)^c * M_{yz}(t)^b) \\ &\quad * (M_{zx}(t)^b * M_{zx}(t)^a * M_{zx}(t)^c) \\ &\geq (M_{xy}(t)^c \cdot M_{xy}(t)^b \cdot M_{xy}(t)^a) * (M_{yz}(t)^a \cdot M_{yz}(t)^c \cdot M_{yz}(t)^b) \\ &\quad * (M_{zx}(t)^b \cdot M_{zx}(t)^a \cdot M_{zx}(t)^c) \\ &= M_{xy}(t)^{a+b+c} * M_{yz}(t)^{a+b+c} * M_{zx}(t)^{a+b+c} \\ &\geq M_{xy}(t) * M_{yz}(t) * M_{zx}(t) = \theta(t). \end{aligned}$$

We find that $\theta(kt) \geq \theta(t)$ implies that $\theta(t) \geq \theta(t/k) \geq \theta(t/k^2) \geq \dots \geq \theta(t/k^n)$ for all $t > 0$ and $n \geq 1$. By (20), (21) and (22),

$$\begin{aligned} M_{xy}(kt) &\geq M_{yz}(t)^a * M_{zx}(t)^b * M_{xy}(t)^c \geq M_{yz}(t) * M_{zx}(t) * M_{xy}(t) = \theta(t) \geq \theta(t/k^n), \\ M_{yz}(kt) &\geq M_{zx}(t)^a * M_{xy}(t)^b * M_{yz}(t)^c \geq M_{zx}(t) * M_{xy}(t) * M_{yz}(t) = \theta(t) \geq \theta(t/k^n), \\ M_{zx}(kt) &\geq M_{xy}(t)^a * M_{yz}(t)^b * M_{zx}(t)^c \geq M_{xy}(t) * M_{yz}(t) * M_{zx}(t) = \theta(t) \geq \theta(t/k^n). \end{aligned}$$

Letting $n \rightarrow \infty$, we have $\lim_{n \rightarrow \infty} \theta(t/k^n) = 1$ for all $t > 0$, and this means that $M_{xy}(kt) = M_{yz}(kt) = M_{zx}(kt) = 1$ for all $t > 0$, i.e., $x = y = z$. The unicity of x follows from (1). \square

Remark 12 The unicity of the coincidence point of F and g is not always true. For instance, if $F \equiv x_0$ is constant and $g \equiv x_0$ is also constant, then every $(x, y, z) \in X^3$ is a coincidence point of F and g .

Remark 13 In the previous theorem, we have only used the continuity of $*$ at $(1, 1)$, that is, if $\{x_n\}, \{y_n\} \subset [0, 1]$ are sequences such that $\{x_n\} \rightarrow 1$ and $\{y_n\} \rightarrow 1$, then $\{x_n * y_n\} \rightarrow 1$. And this is true because $\{x_n * y_n\} \geq \{x_n \cdot y_n\} \rightarrow 1 \cdot 1 = 1$.

Example 14 Consider $(X = \mathbb{R}, M^e)$ as in Example 7. Let $\alpha, \beta > 0$ and $k \in (0, 1)$ be positive real numbers such that $6\alpha \leq \beta k$ (in particular, $\alpha/k \leq \beta/6$). Define $F : \mathbb{R}^3 \rightarrow \mathbb{R}$ and $g : \mathbb{R} \rightarrow \mathbb{R}$ as $F(x, y, z) = \alpha(x - y)$ and $gx = \beta x$ for all $x, y, z \in X$. Clearly, g is continuous, F and g are commuting and $F(\mathbb{R}^3) = \mathbb{R} = g(\mathbb{R})$. We also note that M^e verifies

$$\begin{aligned} M_{FxyzFuvw}^e(kt) &= (e^{-(x-u)+(v-y)})^{-\frac{\alpha}{kt}} \geq (e^{-\frac{2 \max(|x-u|, |v-y|)}{t}})^{\frac{\alpha}{k}} \\ &\geq (e^{-\frac{2 \max(|x-u|, |v-y|)}{t}})^{\frac{\beta}{6}} = (e^{-\frac{\beta}{3t}})^{\max(|x-u|, |v-y|)} = \min(e^{-\frac{\beta|x-u|}{3t}}, e^{-\frac{\beta|v-y|}{3t}}) \\ &\geq \min(e^{-\frac{|\beta x - \beta u|}{3t}}, e^{-\frac{|\beta y - \beta v|}{3t}}, e^{-\frac{|\beta z - \beta w|}{3t}}) \\ &= \min([M_{gxgu}^e(t)]^{1/3}, [M_{gygv}^e(t)]^{1/3}, [M_{gzgw}^e(t)]^{1/3}). \end{aligned}$$

Therefore, applying Theorem 11, we deduce that F and g have a tripled coincidence point.

Consequences

In the proof of the next result, the view of (X, d) as the crisp FMS (X, M^c, \min) is used (see Example 7). This approach allows us to deduce results for metric spaces from the corresponding result in the fuzzy setting. Moreover, Theorem 15 is just a tripled coincidence point result, similar to Berinde-Borcut one, see [18, Theorem 7] and [21, Theorem 4], in a not necessarily partially ordered set.

Theorem 15 Let (X, d) be a complete metric space and let $F : X^3 \rightarrow X$ and $g : X \rightarrow X$ be two mappings such that $F(X^3) \subseteq g(X)$ and g is continuous and commuting with F . Suppose that F and g verify some of the following conditions for all $x, y, z, u, v, w \in X$:

- (a) $d_{FxyzFuvw} \leq k \max(d_{gxgu}, d_{gygv}, d_{gzgw})$ for some $k \in (0, 1)$.
- (b) $d_{FxyzFuvw} \leq k(\alpha d_{gxgu} + \beta d_{gygv} + \gamma d_{gzgw})$ for some $k \in (0, 1)$ and some $\alpha, \beta, \gamma \in [0, 1/3]$.
- (c) $d_{FxyzFuvw} \leq \alpha d_{gxgu} + \beta d_{gygv} + \gamma d_{gzgw}$ for some $\alpha, \beta, \gamma \in [0, 1)$ such that $\alpha + \beta + \gamma < 1$.

Then there exists a unique $x \in X$ such that $x = gx = F_{xxx}$.

Proof (a) Consider M^c defined as in Example 7. As (X, d) is complete, then (X, M^c, \min) is a complete FMS. Fix $x, y, z, u, v, w \in X$ and $t > 0$, and we are going to prove (1) using $a = b = c = 1/3$ and $*$ = min. If $M_{gxgu}^c(t) = 0$ or $M_{gygv}^c(t) = 0$ or $M_{gzgw}^c(t) = 0$, then (1) is obvious. Suppose that $M_{gxgu}^c(t) = 1$, $M_{gygv}^c(t) = 1$ and $M_{gzgw}^c(t) = 1$. This means that $d_{gxgu} < t$, $d_{gygv} < t$ and $d_{gzgw} < t$. Therefore, $t > \max(d_{gxgu}, d_{gygv}, d_{gzgw})$ and $kt > k \max(d_{gxgu}, d_{gygv}, d_{gzgw}) \geq d_{F_{xyz}F_{uvw}}$. Hence, $M_{F_{xyz}F_{uvw}}^c(kt) = 1$ and (1) is also true.

(b) In this case,

$$\begin{aligned} d_{F_{xyz}F_{uvw}} &\leq k(\alpha d_{gxgu} + \beta d_{gygv} + \gamma d_{gzgw}) \leq k\left(\frac{1}{3}d_{gxgu} + \frac{1}{3}d_{gygv} + \frac{1}{3}d_{gzgw}\right) \\ &= \frac{k}{3}(d_{gxgu} + d_{gygv} + d_{gzgw}) \leq \frac{k}{3}3 \max(d_{gxgu}, d_{gygv}, d_{gzgw}) \\ &= k \max(d_{gxgu}, d_{gygv}, d_{gzgw}). \end{aligned}$$

(c) If $k = \alpha + \beta + \gamma < 1$,

$$\begin{aligned} d_{F_{xyz}F_{uvw}} &\leq \alpha d_{gxgu} + \beta d_{gygv} + \gamma d_{gzgw} \leq \alpha \max(d_{gxgu}, d_{gygv}, d_{gzgw}) \\ &\quad + \beta \max(d_{gxgu}, d_{gygv}, d_{gzgw}) + \gamma \max(d_{gxgu}, d_{gygv}, d_{gzgw}) \\ &= (\alpha + \beta + \gamma) \max(d_{gxgu}, d_{gygv}, d_{gzgw}) = k \max(d_{gxgu}, d_{gygv}, d_{gzgw}). \quad \square \end{aligned}$$

Example 16 If $X = \mathbb{R}$, $d(x, y) = |x - y|$ for all $x, y \in \mathbb{R}$ and $a, b, c, d, M \in \mathbb{R}$ are such that $M > |a| + |b| + |c|$, the mappings $F : \mathbb{R}^3 \rightarrow \mathbb{R}$ and $g : \mathbb{R} \rightarrow \mathbb{R}$, defined as $F_{xyz} = (ax + by + cz + d)/M$ and $gx = x$ for all $x, y, z \in \mathbb{R}$, verify the hypothesis of Theorem 15(c). It is easy to check that (x_0, x_0, x_0) , where $x_0 = d/(M - a - b - c)$, is the unique tripled coincidence point of F and g and verifies $F(x_0, x_0, x_0) = x_0$.

Now, we prove the existence of a coupled coincided point for $F : X^2 \rightarrow X$ and g that generalizes Theorem 4.1 in [7], taking $a = b = 1/2$. That is, the main result of the paper also covers the main theoretical results of Zhu and Xiao [7].

Corollary 17 Let $*$ be a t -norm of H -type such that $s * t \geq st$ for all $s, t \in [0, 1]$. Let $k \in (0, 1)$ and $a, b \in [0, 1]$ be real numbers such that $a + b \leq 1$, let $(X, M, *)$ be a complete FMS and let $F : X^2 \rightarrow X$ and $g : X \rightarrow X$ be two mappings such that $F(X^2) \subseteq g(X)$ and g is continuous and commuting with F . Suppose that

$$M_{F_{xy}F_{uv}}(kt) \geq M_{gxgu}(t)^a * M_{gygv}(t)^b$$

for all $x, y, u, v \in X$ and all $t > 0$. Then there exists a unique $x \in X$ such that $x = gx = F_{xx}$.

Proof Define $c = 0$ and $F' : X^3 \rightarrow X$ as $F'_{xyz} = F_{xy}$ for all $x, y, z \in X$. Then $F'(X^3) = F(X^2) \subseteq g(X)$ and F' is commuting with g ($gF'_{xyz} = gF_{xy} = F_{gxy} = F'_{gxgygz}$). Furthermore,

$$\begin{aligned} M_{F'_{xyz}F'_{uvw}}(kt) &= M_{F_{xy}F_{uv}}(kt) \geq M_{gxgu}(t)^a * M_{gygv}(t)^b \\ &= M_{gxgu}(t)^a * M_{gygv}(t)^b * 1 \geq M_{gxgu}(t)^a * M_{gygv}(t)^b * M_{gzgw}(t)^c. \end{aligned}$$

Then there exists a unique $x \in X$ such that $gx = F'_{xxx}$. If $y \in X$ verifies $F_{yy} = gy$, then $gy = F_{yy} = F'_{yyy}$, so $x = y$. □

Corollary 18 ([1, Theorem 2.2]) *Let (X, d) be a complete metric space and let $F : X^2 \rightarrow X$ and $g : X \rightarrow X$ be two mappings such that $F(X^2) \subseteq g(X)$ and g is continuous and commuting with F . Suppose that F and g verify some of the following conditions for all $x, y, u, v \in X$:*

- (a) $d_{F_{xy}F_{uv}} \leq k \max(d_{g_xg_u}, d_{g_yg_v})$ for some $k \in (0, 1)$.
- (b) $d_{F_{xy}F_{uv}} \leq k(\alpha d_{g_xg_u} + \beta d_{g_yg_v})$ for some $k \in (0, 1)$ and some $\alpha, \beta \in [0, 1/2]$.
- (c) $d_{F_{xy}F_{uv}} \leq \alpha d_{g_xg_u} + \beta d_{g_yg_v}$ for some $\alpha, \beta, \gamma \in [0, 1)$ such that $\alpha + \beta < 1$.

Then there exists a unique $x \in X$ such that $x = gx = F_{xx}$.

Proof Similar to the proof of Theorem 15. □

Remark 19 In fact, the previous result is proved for X , a partially ordered set in [1].

Moreover, from a similar procedure, we can deduce the celebrated Banach contraction principle (Theorem 1).

Applications Lipschitzian systems

Let $f_1, f_2, f_3 : \mathbb{R} \rightarrow \mathbb{R}$ be Lipschitzian mappings and let $\beta_1, \beta_2, \beta_3 \in \mathbb{R}$ be real numbers. Define $h : \mathbb{R} \rightarrow \mathbb{R}$ as $h(x) = \beta_1 f_1(x) + \beta_2 f_2(x) + \beta_3 f_3(x)$ for all $x \in \mathbb{R}$. Then h is another Lipschitzian mapping and $k_h \leq |\beta_1|k_{f_1} + |\beta_2|k_{f_2} + |\beta_3|k_{f_3}$. Obviously, if $K = |\beta_1|k_{f_1} + |\beta_2|k_{f_2} + |\beta_3|k_{f_3} < 1$, then h is a contraction, so there exists a unique $x_0 \in \mathbb{R}$ such that $h_{x_0} = x_0$.

Next, define $F : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ as $F_{xyz} = \beta_1 f_1(x) + \beta_2 f_2(y) + \beta_3 f_3(z)$ for all $x, y, z \in \mathbb{R}$. It is clear that $F_{xxx} = h_x$ for all $x \in \mathbb{R}$. Furthermore,

$$d(F_{x_1x_2x_3}, F_{y_1y_2y_3}) \leq \sum_{i=1}^3 |\beta_i| |f_i(x_i) - f_i(y_i)| \leq \sum_{i=1}^3 |\beta_i| k_{f_i} |x_i - y_i| \leq K \max_{1 \leq j \leq 3} d(x_j, y_j).$$

If $K < 1$, then F verifies (1) with $gx = x$ for all $x \in \mathbb{R}$.

Corollary 20 *Let $f_1, f_2, f_3 : \mathbb{R} \rightarrow \mathbb{R}$ be Lipschitzian mappings on \mathbb{R} (provided with the Euclidean metric) and let $\beta_1, \beta_2, \beta_3 \in \mathbb{R}$ such that $|\beta_1|k_{f_1} + |\beta_2|k_{f_2} + |\beta_3|k_{f_3} < 1$. Then the system*

$$(S) \begin{cases} \beta_1 f_1(x) + \beta_2 f_2(y) + \beta_3 f_3(z) = x, \\ \beta_1 f_1(y) + \beta_2 f_2(z) + \beta_3 f_3(x) = y, \\ \beta_1 f_1(z) + \beta_2 f_2(x) + \beta_3 f_3(y) = z \end{cases}$$

has a unique solution, which is (x_0, x_0, x_0) , where x_0 is the only real solution of $\beta_1 f_1(x) + \beta_2 f_2(x) + \beta_3 f_3(x) = x$.

Example 21 Consider the system

$$(S_1) \begin{cases} 30 \sin x - \frac{28}{1+y^2} + 150 = 72x - 15 \arctan z, \\ 30 \sin y - \frac{28}{1+z^2} + 150 = 72y - 15 \arctan x, \\ 30 \sin z - \frac{28}{1+x^2} + 150 = 72z - 15 \arctan y. \end{cases}$$

If we choose $f_1(x) = 5 + \sin x$, $f_2(x) = 1/(1 + x^2)$ and $f_3(x) = \arctan x$, then f_1, f_2 and f_3 are Lipschitzian mappings, and $k_{f_1} = k_{f_3} = 1$ and $k_{f_2} = 3\sqrt{3}/8$. Let $\beta_1 = 5/12$, $\beta_2 = -7/18$ and $\beta_3 = 5/24$. Then $|\beta_1|k_{f_1} + |\beta_2|k_{f_2} + |\beta_3|k_{f_3} = (30 + 7\sqrt{3})/48 < 1$. As system (S_1) is equal to (S) , then (S_1) has a unique solution, which is of the form (x_0, x_0, x_0) , where x_0 is the only solution of

$$30 \sin x - \frac{28}{1 + x^2} + 150 = 72x - 15 \arctan x.$$

Finding, for example, the root by the bisection method, we get, approximately, $x_0 = 2.5212648363927$.

An integral system

Let $a, b \in \mathbb{R}$ with $a < b$ and let $I = [a, b]$. Consider $X = \mathcal{L}^1(I)$ with the distance $d_1(f, g) = \int_I |f(t) - g(t)| dt$, where \int represents the Lebesgue integral. It is well known that $(\mathcal{L}^1(I), d_1)$ is a complete MS. Let $k, \beta_1, \beta_2, \beta_3 \in \mathbb{R}$ be real numbers and let $G : \mathbb{R}^3 \rightarrow \mathbb{R}$ be a mapping verifying $G(0, 0, 0) = 0$ and

$$|G_{x_1 x_2 x_3} - G_{y_1 y_2 y_3}| \leq k \sum_{i=1}^3 \beta_i |x_i - y_i| \quad \text{for all } (x_1, x_2, x_3), (y_1, y_2, y_3) \in \mathbb{R}^3.$$

If $A \in \mathbb{R}$, we want to find functions $f_1, f_2, f_3 \in \mathcal{L}^1(I)$ such that

$$f_i(x) = A + \int_{[a,x]} G(f_i(t), f_{i+1}(t), f_{i+2}(t)) dt \tag{23}$$

holds for all $x \in I, i = 1, 2, 3$.

For all $f_1, f_2, f_3 \in \mathcal{L}^1(I)$ and all $x \in I$, define

$$F_{f_1 f_2 f_3}(x) = A + \int_{[a,x]} G(f_1(t), f_2(t), f_3(t)) dt.$$

On the one hand, it is not difficult to prove that $F_{f_1 f_2 f_3} \in \mathcal{L}^1(I)$, hence $F : \mathcal{L}^1(I)^3 \rightarrow \mathcal{L}^1(I)$ is well defined. On the other hand,

$$\begin{aligned} d_1(F_{f_1 f_2 f_3}, F_{g_1 g_2 g_3}) &= \int_I |F_{f_1 f_2 f_3}(x) - F_{g_1 g_2 g_3}(x)| dx \\ &\leq \int_I \left(\int_{[a,x]} |G(f_1(t), f_2(t), f_3(t)) - G(g_1(t), g_2(t), g_3(t))| dt \right) dx \\ &\leq \int_I \left(\int_{[a,x]} k \sum_{i=1}^3 \beta_i |f_i(t) - g_i(t)| dt \right) dx \\ &\leq k \sum_{i=1}^3 \beta_i \int_I \left(\int_I |f_i(t) - g_i(t)| dt \right) dx \\ &= k \sum_{i=1}^3 \beta_i \int_I d_1(f_i, g_i) dx = k(b-a) \sum_{i=1}^3 \beta_i d_1(f_i, g_i). \end{aligned}$$

If we suppose that $K = k(b-a)(\beta_1 + \beta_2 + \beta_3) < 1$, then F verifies (1) with $g(f) = f$ for all $f \in \mathcal{L}^1(I)$. Then the system (23) has a unique solution, which is of the form (f_0, f_0, f_0) , where

$f_0 \in \mathcal{L}^1(I)$ is the only solution of the equation

$$f_0(x) = A + \int_{[a,x]} G(f_0(t), f_0(t), f_0(t)) dt \quad \text{for all } x \in I$$

(this exists as a simple application of the *Banach contraction principle*).

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors completed the paper together. All authors read and approved the final manuscript.

Author details

¹Department of Statistics and Operations Research, University of Jaén, Jaén, Spain. ²Department of Mathematics, University of Jaén, Jaén, Spain. ³Department of Statistics and Operations Research, University of Granada, Granada, Spain.

Received: 1 July 2012 Accepted: 11 January 2013 Published: 11 February 2013

References

1. Bhaskar, T, Lakshmikantham, V: Fixed point theorems in partially ordered metric spaces and applications. *Nonlinear Anal. TMA* **65**(7), 1379-1393 (2006)
2. Lakshmikantham, V, Ćirić, L: Coupled fixed point theorems for nonlinear contractions in partially ordered metric spaces. *Nonlinear Anal. TMA* **70**, 4341-4349 (2009)
3. Fang, J: Common fixed point theorems of compatible and weakly compatible maps in Menger spaces. *Nonlinear Anal. TMA* **5-6**, 1833-1843 (2009)
4. Shakeri, S, Ćirić, L, Saadati, R: Common fixed point theorem in partially ordered L -fuzzy metric spaces. *Fixed Point Theory Appl.* **2010**, Article ID 125082 (2010)
5. Sedghi, S, Altun, I, Shobe, N: Coupled fixed point theorems for contractions in fuzzy metric spaces. *Nonlinear Anal. TMA* **72**(3-4), 1298-1304 (2010)
6. Hu, X: Common coupled fixed point theorems for contractive mappings in fuzzy metric spaces. *Fixed Point Theory Appl.* **2011**, Article ID 363716 (2011)
7. Zhu, X, Xiao, J: Note on "Coupled fixed point theorems for contractions in fuzzy metric spaces". *Nonlinear Anal. TMA* **74**(16), 5475-5479 (2011)
8. Sintunavarat, W, Cho, Y, Kumam, P: Coupled coincidence point theorems for contractions without commutative condition in intuitionistic fuzzy normed spaces. *Fixed Point Theory Appl.* **2011**, 81 (2011)
9. Zadeh, L: Fuzzy sets. *Inf. Control* **8**, 338-353 (1965)
10. George, A, Veeramani, P: On some results in fuzzy metric spaces. *Fuzzy Sets Syst.* **64**, 395-399 (1994)
11. Kramosil, I, Michalek, J: Fuzzy metric and statistical metric spaces. *Kybernetika* **11**, 326-333 (1975)
12. Grabiec, M: Fixed points in fuzzy metric spaces. *Fuzzy Sets Syst.* **27**(3), 385-389 (1988)
13. Fang, J: On fixed point theorems in fuzzy metric spaces. *Fuzzy Sets Syst.* **46**, 107-113 (1992)
14. Cho, Y: Fixed points in fuzzy metric spaces. *J. Fuzzy Math.* **5**(4), 949-962 (1997)
15. Gregori, V, Sapena, A: On fixed-point theorems in fuzzy metric spaces. *Fuzzy Sets Syst.* **125**(2), 245-252 (2002)
16. Beg, I, Abbas, M: Common fixed points of Banach operator pair on fuzzy normed spaces. *Fixed Point Theory* **12**(2), 285-292 (2011)
17. Elagan, SK, Rahmat, MS: Some fixed points theorems in locally convex topology generated by fuzzy n -normed spaces. *Iran. J. Fuzzy Syst.* **9**(4), 43-54 (2012)
18. Berinde, V, Borcut, M: Tripled fixed point theorems for contractive type mappings in partially ordered metric spaces. *Nonlinear Anal. TMA* **74**, 4889-4897 (2011)
19. Roldán, A, Martínez-Moreno, J, Roldán, C: Multidimensional fixed point theorems in partially ordered metric spaces. *J. Math. Anal. Appl.* **396**, 536-545 (2012)
20. Hadžić, O, Pap, E: *Fixed Point Theory in Probabilistic Metric Spaces*. Kluwer Academic, Dordrecht (2001)
21. Borcut, M, Berinde, V: Tripled coincidence theorems for contractive type mappings in partially ordered metric spaces. *Appl. Math. Comput.* **218**(10), 5929-5936 (2012)

doi:10.1186/1687-1812-2013-29

Cite this article as: Roldán et al.: Tripled fixed point theorem in fuzzy metric spaces and applications. *Fixed Point Theory and Applications* 2013 **2013**:29.